

**CLIMATIC CHANGES:  
Anthropogenic Influence  
Or  
Naturally Induced Phenomenon**

**By**

**Anthony Foscolos**

# **CLIMATIC CHANGES: Anthropogenic Influence or Naturally Induced Phenomenon**

**By**

**A.E. Foscolos**

**Department of Mineral Resources Engineering  
Technical University of Crete**

By the end of the 18<sup>th</sup> century eminent scientists (Franklin, 1784; Fourier, 1824; 1827; Agassiz, 1840; Tyndal, 1859; Croll, 1864; Koppen, 1873; Czerney, 1881; Arrhenius, 1896) explained the climatic changes on the basis of temperature and the ensuing glacial retreat. This disturbing observation led many prominent scientists (de Saussure, Bunsen, Pettenkoffer, Kroch<sup>1</sup> and Warburg<sup>1</sup>) to send air balloons equipped with special devices to trap air from the lower atmosphere in order to measure CO<sub>2</sub> concentrations. Ninety thousand (90,000) measurements were carried out at 138 locations in 4 continents between 1810 and 1961.

The data indicated that atmospheric CO<sub>2</sub> concentrations, during the 19<sup>th</sup> century varied between 290 ppm and 430 ppm (with an average of 322 ppm for the pre-industrial period). For the 20<sup>th</sup> century, the average concentration is 338 ppm when combined with comparable CO<sub>2</sub> measurements carried out by Mauna Loa Observatory, Hawaii, USA, (1958- 2000). Measurement precision is +/- 3%.

Based on thermometric measurements, the mean average temperature increase from 1850 to the present is 0.75°C (0.44°C/100 years) with the following fluctuations. From 1850 to 1940 the temperature increased by +0.6°C; while from 1941 to 1975 temperature dropped by -0.2°C. From 1976 to 1998, the temperature rose by +0.35°C. From 1999 to 2006 temperature increase was nil. Finally, since 2007 the Mean Annual Temperature of earth's surface has substantially decreased.

As far as CO<sub>2</sub> concentration in the air's atmosphere is concerned, it has been well documented that during the Holocene Epoch there is a substantial

---

<sup>1</sup> Nobel Prize Winners in Science, 1920 and 1931, respectively

time lag between maximum temperatures recorded during the Interglacial periods and maximum CO<sub>2</sub> concentrations in the atmosphere. Moreover, the same time lag is documented between 1850 and 1980, where CO<sub>2</sub> concentrations in the atmosphere lag behind the increase of temperature for more than 100 years. A parallel increase of CO<sub>2</sub> concentrations in the atmosphere and temperature increase is observed only between 1981 and 1995. No correlation is seen thereafter. The divergence is substantial from 2003 to the present where CO<sub>2</sub> concentrations are increased while temperatures are decreased.

These interpretations exclude any correlation between atmospheric CO<sub>2</sub> concentrations and temperature fluctuations. Hence, in order to explain the well documented climatic changes the influence of many natural climate drivers should be accepted.

**Key words:** Climatic changes, temperatures, CO<sub>2</sub> concentrations

## **Contents**

- 1. Introduction – Brief Historical Review**
- 2. Analysis of existing climatic changes data**
  - a) Middle Pleistocene to Holocene*
  - b) Holocene (Last Interglacial period)*
    - i. 12000 BC to 1850 AD*
    - ii. 1850 AD to 2008 AD*
- 3. Measurements of atmospheric CO<sub>2</sub> concentrations**
- 4. The assumed correlation between atmospheric CO<sub>2</sub> and temperature**
- 5. Relation of Temperature to Solar activity**
- 6. Other Climatic Anomalies**
- 7. Conclusions**
- 8. References**

# 1. Introduction

## Brief Historical Review

Climatic changes have been the subject of intensive research since the late 18<sup>th</sup> century by eminent scientists (Franklin, 1784; Fourier, 1824; 1827; Agassiz, 1840; Tyndal, 1861, 1863; Croll, 1864; Koppen, 1873; Czerney, 1881; Arrhenius, 1896, 1905). They developed theories to link the presence of erratic boulders in various places in the world to action of former glaciers as well as to explain the temperature rise.

In addition to the various theories and observations, air balloons equipped with special devices to trap air from the lower atmosphere were sent from a number of European scientists (de Saussure, Bunsen, Pettenkoffer, Kroch and Warburg) in order to measure CO<sub>2</sub> concentrations (Beck 2007). Ninety thousand (90,000) measurements were carried out at 138 locations in 4 continents between 1810 and 1961. The data indicated that atmospheric CO<sub>2</sub> concentrations varied between 290 ppm and 430 ppm during the 19<sup>th</sup> century (with an average for the pre-industrial period of 322 ppm), Figure 1. For the 20<sup>th</sup> century, the average concentration is 338 ppm when combined with comparable CO<sub>2</sub> measurements carried out by Mauna Loa Observatory (1958- 2009) (Atmospheric CO<sub>2</sub> at Mauna Loa Observatory, Hawaii, USA, 2009). Measurement precision were of the order of +/- 3%.




Figure 1. Evidence of variability of atmospheric CO<sub>2</sub> concentration during the 20<sup>th</sup> century in the Northern Hemisphere Beck, 2007.

In addition to this intense research work, field geologists have mapped the Quaternary glaciations extent and their chronology in Europe as well as in North and South America. Altogether, it took several decades until the ice age theory was fully accepted. This happened on an international scale in the second half of the 1870's (Kruger, 2008). This work is summarized in Figure 2 and Table 1.




Figure 2. Late Pleistocene Epoch: Atmospheric CO<sub>2</sub> and the Glacial Cycles. Time Line Glaciations, 2009. Pleistocene Climate Cycles. [www.en.wikipedia.org/wiki/Timeline\\_of\\_glaciation](http://www.en.wikipedia.org/wiki/Timeline_of_glaciation).

**Table 1. Land-based chronology of Pleistocene glacial cycles. Time Line Glaciations, 2009. Pleistocene Climate Cycles.**  
[www.en.wikipedia.org/wiki/Timeline\\_of\\_glaciation](http://www.en.wikipedia.org/wiki/Timeline_of_glaciation).

Backwards Glacial Index	Names						Inter/ Glacial Period (ka)	Epoch	
	Alpine	N. American	N. European	Great Britain	S. American				
				<a href="#">Flandrian</a>			interglacial	present – 12	<a href="#">Holocene</a>
1 <sup>st</sup>	<a href="#">Würm</a>	<a href="#">Wisconsin</a>	<a href="#">Weichsel</a> or Vistula	<a href="#">Devensian</a>	<a href="#">Llanquihue</a>		glacial period	12 – 110	
	<a href="#">Riss- Würm</a>	<a href="#">Sangamonian</a>	<a href="#">Eemian</a>	<a href="#">Ipswichian</a>			interglacial	110 – 130	
2 <sup>nd</sup>	<a href="#">Riss</a>	<a href="#">Illinoian</a>	<a href="#">Saale</a>	<a href="#">Wolstonian</a> or Gipping	<a href="#">Santa María</a>		glacial period	130 – 200	
	<a href="#">Mindel- Riss</a>	<a href="#">Yarmouth</a>	<a href="#">Holstein</a>	<a href="#">Hoxnian</a>			interglacial(s)	200 – 300/380	<a href="#">Pleistocene</a>
3 <sup>rd</sup> – 5 <sup>th</sup>	<a href="#">Mindel</a>	<a href="#">Kansan</a>	<a href="#">Elsterian</a>	<a href="#">Anglian</a>	Río Llico		glacial period(s)	300/380 – 455	
	<a href="#">Günz- Mindel</a>	<a href="#">Aftonian</a>		<a href="#">Cromerian</a> *			interglacial(s)	455 – 620	
7 <sup>th</sup>	<a href="#">Günz</a>	<a href="#">Nebraskan</a>	<a href="#">Menapian</a>	<a href="#">Beestonian</a>	Caracol		glacial period	620 – 680	


## 2. Analysis of existing climatic changes data

### a) Middle Pleistocene to Holocene Epochs


Recent ice coring data from Vostok-1 (Petit et. al., 1997 and 1999) and EPICA (European Project for Ice Coring in Antarctica, Epica 2004) not only have concurred about the climatic changes but also show interglacial temperatures of 1°C to 2.5°C higher than the present Mean Annual Temperature of the Northern Hemisphere of 15°C, for thousands of years Figure 3. Since proxy temperature measurements of  $\delta^{18}\text{O}$ , were carried out on ice core samples it means that even at temperatures well above those prevailing today's, polar ice caps did not melt. These large climatic changes were theoretically attributed to the eccentricity, tilting and wobbling of the earth (Milankovitch, 1940). Finally after three decades this theory was scientifically accepted (Hayes et. al., 1974).

Geomagnetic polarity technique along with appropriate sampling has provided a direct assessment of glacial and interglacial environments (Barendregt and Duk-Rodkin, 2004). Through these studies, and  $\delta^{18}\text{O}$  paleotemperature record from Site 607 in North America, over 100 glacial and interglacial periods were identified (Ruddiman et al., 1989; Raymo, 1992) (Figure 4). Therefore, it is

seen that from the Pliocene till the Holocene Epoch, climatic changes were the norm.



**Figure 3.** Climatic changes as documented from Vostok-1 icecore data, Petit et. al., 1999, and EPICA icecore data, EPICA, 2004, for the last 450,000 years. Worth noticing is the rise of temperature well above the today's one during the long interglacial periods without the complete melting of the ice caps



**Figure 4. Geomagnetic polarity scale (Candle and Kent 1995) and  $\delta^{18}\text{O}$  paleotemperatures record from site 607 in the North Atlantic (Ruddiman et.al., 1989; Raymo 1992)**



**b) Holocene Epoch. Last Interglacial period:**

**i. 12000 BC to 1850 AD**

Climatic changes during this period are very well documented by Dansgaard et al., 1969 and Schonwiese, 1995, Figure 5. The recent glacial retreat began about 14000 years ago (12000 BC). This warming period was shortly interrupted by a sudden cooling, known as Younger-Dryas, at about 10000 to 8500 BC. From 8000 BC to about 4000 BC the average global temperature reached its maximum level during the Holocene Epoch and was 1°C to 2°C warmer than today's Mean Annual Temperature of the Earth's Atmosphere of 15°C (Pidwirny, 2006). Climatologists call this period the Climatic Optimum. Worth noticing is that polar ice caps did not vanish between 8000 BC to 4000 BC, since ice core samples were recently collected and CO<sub>2</sub> concentrations were measured in air bubbles which were trapped in the ice cores. Between the span of 4000 years, 2 minor cooling events took place, while a substantial cooling trend took place between 3500 BC and 2000 BC.





Figure 5. Average near surface temperatures of the northern hemisphere during the past 11000 years Dansgaard et al., 1969 & Schonwiese 1995.

From 450 BC to 150 AD, Northern Europe was subjected to another warm period the so-called Roman Warm Period with average temperatures of 2.5°C higher than today's temperature ( Keigwin, 1996; Holmgren et al., 1999; 2001; Idso and Idso, 2000; Olafsdottir et al., 2001; Grudd et al., 2002; Jiang et al., 2002; Berglund, 2003; Munroe 2003; D' Arrigo et al., 2004; Loehle, 2004; Fleitman et al., 2004; Hormes et al., 2004; Blundel and Barber, 2005; Linderholm and Gunnarson, 2005; Allen et al., 2007; Mariolakos, 2008) (Figure 6). Again worth noticing is the fact that polar ice caps did not vanish during the 600 years

time span since, again, temperatures were measured in ice core samples and CO<sub>2</sub> concentrations were measured in air bubbles trapped within the ice cores, as well. Subsequently, a cooling period has begun; the last one was until about 900 AD. At its height, the cooling caused the Nile River and the Black Sea to freeze, 829 AD and 800 AD to 801 AD respectively, (Pidwirny, 2006).



**Figure 6. Sea surface temperatures from the Sargasso sea during the last 3000 years based on oxygen isotopic ratios of Globigerinoids rubber (Plankton) collected from a box core through 50 cm of bottom sediment, Keigwin, 1996**

The period from 900 to 1350 is called the Medieval Warm Period (MWP). During this period, temperatures fluctuated from +0.4°C above the today's (Soon and Baliunas, 2003; Moberg et al., 2005; Viau et al., 2006; Loehle 2007; Loehle and Mc Culloch, 2008), (Figure 7), to +0.8°C higher than today's (Seppa and Birks, 2001, 2002; Heikkila and Seppa, 2003). Their estimate was based on pollen data in order to reconstruct past climate thus studying Fennoscandian tree-line fluctuations. The existence of Medieval Warm Period was challenged by Mann et al. (1998; 1999) (Figure 8) based upon proxy measurements of temperatures from the width of tree rings. However, tree ring data may not capture long-term climate changes (100+ years) because tree size, root/shoot ratio, genetic adaptation to climate, and forest density can all shift in response to prolonged climate changes, among other reasons (Broecker, 2001; Falcon-Lang, 2005; Loehle, 2005; Moberg et al., 2005).




Figure 7. 2000 years of global temperatures based upon 30 year averages, Spencer, 2007.




Figure 8. Millennial Northern Hemisphere (NH) Temperature Reconstruction (blue line) and Instrumental Data (red line from 1000 to 1999 AD, Mann et. al., 1999). The graph, “Hockey Stick” relies mainly on tree-rings studies which are annual cycles (high frequency). It is known from geophysics that it is difficult to obtain low frequency (centennial change) data where low frequency is filtered. This graph denies historical records of Medieval Warm Period and Little Ice Age well described by historians, Leroy-Ladurie, 1988 and Lamb, 1995.

Most seriously, typical reconstructions assume that tree ring width is linearly related to temperature, but trees may be related in an inverse parabolic manner to temperature, with ring width rising with temperature to an optimal level and then decreasing with further temperature increase (D'Arrigo et al., 2004; Kelly et al., 1994). This response is most likely due to water limitation at higher temperatures due to increase of the evaporation rates. The result of this violation of linearity is to introduce tremendous uncertainty or bias into any reconstruction, particularly for temperatures outside the calibration range. For example, tree rings in many places show recent divergence from observed warming trends, even showing downward trends (Briffa et al., 1998a, 1998b; Pisaric et al., 2007). Other important facts that support the existence of a Medieval Warm Period are archaeological and agricultural data. It is well documented that Greenland was settled from 900 AD till 1350 AD and farming took place due to milder weather than today's (Brown, 2000).

Also the mere fact that vineyards were extended to North and South England (Schmidt, 2006) during this period indicates, beyond any shadow of doubt, that the climate was much warmer at that time by at least  $+0.4^{\circ}\text{C}$ . To have vineyards extending 450 km north of their present northerly growing limit, the climate had to be similar to the one we have today in Southern California, Greece, Italy, Spain etc. Hence, the Hockey Stick diagram (Mann's et al., 1998) showing that the  $+0.4^{\circ}\text{C}$  during the MWP is well below the acceptable Mean Annual Temperature of  $15^{\circ}\text{C}$  and more or less equivalent to the Little Age Temperatures while the today's  $+0.35^{\circ}\text{C}$  is well above the acceptable Mean Annual Temperature of  $15^{\circ}\text{C}$ , Figure 8, is completely unacceptable. Also unacceptable is the correlation of the proxy measurements of temperatures based on tree rings, 1000 AD to 1850 AD, with those obtained from thermometers, 1850 AD to 2008 AD.

From the above it is concluded that climatic changes were taking place in the past regardless the presence or absence of human beings on earth, irrespective of atmospheric  $\text{CO}_2$  concentrations and without the use of hydrocarbons. An excellent scientific review of thousands of papers along with new scientific data which refutes the alarmist mantra is presented by Singer and Idso, 2009.

#### *ii. 1850 AD to 2008 AD*

From 1850 thermometric data indicating the Average Mean Annual Temperature of Earth's Surface become available (Jones, 2008; Hadley Meteorolo-

logical Station, 2008) (Figure 9). From 1850 to 1910 temperatures were more or less stable fluctuating between  $-0.4^{\circ}\text{C}$  and  $-0.6^{\circ}\text{C}$  below the optimum average of  $15^{\circ}\text{C}$ . Since 1910, temperatures increased by  $+0.6^{\circ}\text{C}$  to reach the optimum of  $15^{\circ}\text{C}$  in 1940. According to NASA's newly published data the hottest year on record in the USA is 1934 and not 1998; three of the hottest years on record occurred before 1940 and six of the top 10 hottest years occurred before 1960<sup>2</sup>. Afterwards temperatures dropped by  $-0.2^{\circ}\text{C}$  and stayed at this level, from 1940 to 1980. Subsequently, from 1980 to 1998 the Mean Annual Temperature increased by  $+0.35^{\circ}\text{C}$  and remained at this level till 2007. From that date till January 2008 temperature dropped by  $-0.1^{\circ}\text{C}$ , Figure 9. One wonders if this temperature increase, few tenths of a degree, from either 1940 or from 1980 can be considered alarming when in the past much higher temperatures were noticed for extremely long periods without any real damage to either earth or its inhabitants.




Figure 9. Global temperature record 1850-2008, Jones, 2008, and <http://hadobs.metoffice.com/hadcrut3/diagnostics/global/nh+sh/>

Recently, Schneider et. al., (2006), (Figure 10) a series of data suggested that temperatures in Antarctica were colder near the end of the 20<sup>th</sup> century than it was in the early decades of the 19<sup>th</sup> century when atmospheric  $\text{CO}_2$  concentration was about 100 ppm less than it is currently. This is in agreement

<sup>2</sup> <http://data.giss.nasa.gov/gistemp/graphs/ Fig.D.lrg.gif>

with a number of other analyses of Antarctic instrumental surface and air temperature data which also indicate the continent has recently experienced a net cooling, which likely began as early as the mid-1960s (Comiso, 2000; Doran *et al.*, 2002; Thompson and Solomon, 2002)




Figure 10. A 200 year long Antarctic temperature reconstruction (dark line) based upon 200 years sub-annually resolved  $\delta^{18}\text{O}$  and  $\delta\text{D}$  records from precisely dated ice cores obtained from Low Dome Station, Siple Station, Droning Maud Land Station and 2 West Antarctica Sites of the US component of the International Trans-Atlantic Scientific Expedition vs. Mean Temperature of the Southern Hemisphere (lighter line) Schneider *et al.* (2006).

### 3. Measurements of atmospheric $\text{CO}_2$ concentrations

#### *Proxy measurements*

Proxy measurements of atmospheric  $\text{CO}_2$  are carried out using two methods.

The first one relies on the relation between the density of stomata in leaves and atmospheric  $\text{CO}_2$  concentrations. Paleobotanists can take fossilized leaves and count the number of stomata and therefore get a fairly good picture of how much carbon dioxide was in the atmosphere at the time the leaves died (Kurshner *et al.*, 1996; Beerling *et al.*, 1998; Wagner *et al.*, 1999; 2004; Kouwenberg *et al.*, 2003; Kouwenberg, 2004; Kouwenberg *et al.*, 2005), (Figures 11 and 12)




Figure 11. Correlation of stomata frequency ( stomata index=SI) to atmospheric CO<sub>2</sub> from fossil leaves of *B. Pendula* (black circles) and *P. Pubescens* ( white circles) in lake Little Grisbe, Danemark, Wagner et. al, 2004.




Figure 12. Reconstruction of Holocene atmospheric CO<sub>2</sub> values from 8700 BC to 6800 BC based upon measurements a) in air bubbles inclusions from Taylor Dome ice core samples ([www.ngdc.noaa.gov/paleo/taylor.html](http://www.ngdc.noaa.gov/paleo/taylor.html)) and b) stomata indices (SI) from fossilized leaves of *B. Pendula* and *B. Pubescens*, Wagner, et. al., 2004


The second one assumes that, over time, the concentrations of the various atmospheric gasses are locked when the air bubble is "trapped" in ice. And therefore, as long as it can be determined when the air bubble was trapped, the concentration of CO<sub>2</sub> therein and state can be measured, with confidence, that the atmosphere itself had that same concentration at the time the air bubble was trapped.

However there are two wrong assumptions. Ice, though composed mainly of solid water, does still have some molecules that are in a liquid state. Whether a given molecule is in a solid or liquid (or even gaseous) state, at a given time, depends on how much energy that molecule has at that time. As a result, even at low temperatures, among the three main components of the atmosphere, carbon dioxide is seventy (70) times more soluble than nitrogen and thirty (30) times more soluble than oxygen. This means that, when an air bubble is trapped in ice, not only does the liquid in the ice continue to absorb gasses, but it does so selectively, favouring carbon dioxide, by a huge margin, over the other common gasses in the air bubble. Of course, every molecule of carbon dioxide that passes into a solution is removed from the air within the air bubble.

And therefore, since more carbon dioxide is removed, less carbon dioxide will appear in the remaining air. After thousands or millions of years, when that air bubble is tapped, and the gasses within it measured, the concentrations of the various gasses can no longer said to be the same as when that air bubble was trapped, all those years ago (Wiki, Answers, 2009).


Another problem with air bubbles is their dating with respect to the age of ice where trapped in. The consolidation of snow to ice necessary to trap the air takes place at a certain depth (the 'trapping depth') once the pressure of overlying snow is great enough. Since air can freely diffuse from the overlying atmosphere throughout the upper unconsolidated layer (the 'firn'), trapped air is younger than the ice surrounding it. Trapping depth varies with climatic conditions, so the air-ice age difference could vary between 2500 and 6000 years (Barnola et al., 1983, 1987, 1991). This has been acknowledged even by the IPCC scientists by transposing the proxy CO<sub>2</sub> measurements from 1809 to 1892, to read concentrations from 1892 to 1975, (Neftel et. al., 1985; Friedli et. al., 1986), Figure 13. The transposed time was 83 years.





**Figure 13. Concentration of CO<sub>2</sub> in air bubbles from Siple ice core samples, Antarctica, (open squares) and in the atmosphere, 1958-1986, Mauna Loa Observatory (solid line). In a) the original Siple data are given assuming an 83 years younger age of the air bubbles in respect to enclosing ice. In b) The same data after arbitrary shifting/correction of the air bubble age, Neftel et. al., 1985, Friedli et. al., 1986.**

This raises the issue of how the pre-industrial background concentration of CO<sub>2</sub> at 280 ppm has been established (Callendar, 1958). Figure 13, indicates that CO<sub>2</sub> concentration in the atmosphere was not 292 ppm, as stated, but 335 ppm if all the data are considered (Slocum, 1955; Jaworowski et al., 1992). The same background concentration (332 ppm) for the pre-industrial period, is reported by Beck (2007) and Rutledge (2007). Calendar was prejudice in selecting from all his data roughly 30%, which showed concentration around 290 ppm, leaving the remaining 70% which showed concentrations over 300 ppm, Figure 14.



**Figure 14. Average atmospheric CO<sub>2</sub> concentrations measured in the 19<sup>th</sup> and 20<sup>th</sup> centuries, Calendar 1958. Calendar rejected both higher and lower values to arrive at the desired background CO<sub>2</sub> value of 290 ppmv. If he had considered all CO<sub>2</sub> values the concentrations would have been 320 ppmv for the 19<sup>th</sup> century, Jaworowski et. al., 1992, Slocum, 1995.**

It must be noted that the mixing of data from ice-core measurements with the direct and actual atmospheric measurements are questionable because the data obtained from the ice-core measurements are unreliable and they do not reflect paleoatmospheric CO<sub>2</sub> concentrations.

Another inconsistency is the past relation between CO<sub>2</sub> concentrations and temperatures based on the today's existing and very accurate data. Currently, atmospheric CO<sub>2</sub> concentration measurements from Mauna Loa Observatory indicate that CO<sub>2</sub> concentrations increased from 315 ppm to 385 ppm (70ppm) from 1958 to 2008 (Figures 15, 16). From 1940 to date, temperatures, as reported by Jones 2008, increased by +0.35°C above the optimum temperature of 15°C, Figure 9. One wonders, therefore, if CO<sub>2</sub> concentrations in the atmosphere are the driving force behind temperature increase, then why is it that when paleotemperatures, at all interglacial periods, are reported at well over +1°C above the optimum temperature of 15°C, paleo CO<sub>2</sub> concentrations are only 285 ppm. They should have been 460 ppm. Or even higher, if paleotemperatures were +2°C, such as during the Eemian time. The same holds true for the Holocene Epoch period between 8000 AD and 4000 AD when temperatures were +1°C above the optimum temperature of 15°C and the paleoatmospheric CO<sub>2</sub> concentrations were 260 ppm. Henceforth, CO<sub>2</sub> paleoatmospheric concentrations obtained from bubbles yield lower values than stomata indices by 20%,

Figure 12, or by 50% when compared to actual CO<sub>2</sub> measurement concentrations in the atmosphere, Figure 1.




Figure 15. Atmospheric CO<sub>2</sub> concentration from 1958, 315 ppmv, to 2008, 385 ppmv, Atmospheric CO<sub>2</sub> at Mauna Loa Observatory, Hawaii, USA, 2009






Figure 16. Atmospheric CO<sub>2</sub> increase during the last 50 years expressed as a percent of the total air composition. It is the, non discernible, blue line in the bottom, Spencer, 2007  
[www.jbs.org/jbs-news-feed/4333-fifty-years-of-hot-air-](http://www.jbs.org/jbs-news-feed/4333-fifty-years-of-hot-air-)

When comparing the 2 proxy methods it seems that stomata indices (SI), are more reliable as paleoatmospheric CO<sub>2</sub> indicators.

#### ***4. The assumed correlation between atmospheric CO<sub>2</sub> and temperature***

Examining the possible relationship between CO<sub>2</sub> concentration in the atmosphere, temperatures and presence of Polar ice throughout the Phanerozoic Eon, Figure 17, it is obvious that such a relation does not exist. Polar ice existed from the end of the Silurian Period until the beginning of the Ordovician Period that is for 50 million years, while temperatures were low and the atmospheric CO<sub>2</sub> concentration was close to 4000 ppm, that is 10 times higher than today's, Figure 18. Examining the relation between atmospheric CO<sub>2</sub> concentrations and temperatures during the Quaternary Period de-glaciations periods, it is observed that CO<sub>2</sub> increases lags behind temperature increases by 600 +/- 400 years (Khilyuk and Chillingar 2003; 2006), Figure 19. The same is also reported by Fisher et al., (1999), Caillon et al., (2003) and Siegenthaler et. al., (2005). So it seems that atmospheric CO<sub>2</sub> concentrations are the result of temperature increase rather than the driving force. Exactly the same behaviour is observed between 1850 and 1985 where the increase of atmospheric CO<sub>2</sub> concentration lags behind temperature increase (Friis – Christensen and Lassen 1991), Figure 20.



**Figure 17. A. Temperature fluctuation during the Phanerozoic Eon: 500 million years of climate change.**

[en.wikipedia.org/wiki/Geologic\\_temperature\\_record](http://en.wikipedia.org/wiki/Geologic_temperature_record)

**B. Phanerozoic Carbon Dioxide, Royer et. al., 2004**




Figure 18. Correlation of temperature and atmospheric CO<sub>2</sub> concentration over the Phanerozoic time, Berner and Kothavala., 2001, Scotese , 2002




Figure 19. Time lag between maximum temperatures and atmospheric CO<sub>2</sub> concentration during the Quaternary, Khilyuk and Chilingar, 2003.




Figure 20. Correlation between sunspot cycle length, temperature anomalies and atmospheric CO<sub>2</sub> concentration, Friss-Christiensen and Lassen , 1991

Between 1985 and 2000, the atmospheric CO<sub>2</sub> increased along with the temperature which was increased by +0.35°C. This geologically infinitesimal time period, has been used to forecast catastrophic events for mother earth by relating the CO<sub>2</sub> increase in the atmosphere to the rise in temperature. What was omitted was the parallel increase of sunspot numbers which, as will be discussed later, induces temperature increase. However, since 1999, the temperature remained stable until 2006 while CO<sub>2</sub> concentrations increased substantially by 57 billion tons. From 2007 to date, the temperature has been dropping while CO<sub>2</sub> concentrations have risen by an additional 32 billion tons, Figure 21. An increase of 1 ppmv in the atmosphere requires 2.12 Gt of C. One Gt of C requires 3.667 Gt of CO<sub>2</sub>.






Figure 21. World temperature is falling while atmospheric CO<sub>2</sub> is rising. Data for a) Average Mean Temperature of Earth's Surface: UK's Hadley Climate Research Unit CRUT3. b) Lower Troposphere Temperature Measurements: NASA's Microwave Sounding Unit (MSU). c) Atmospheric CO<sub>2</sub> concentration: Mauna Loa Observatory, Hawaii, USA. <http://icecap.us/images/uploads/Correlation Last Decade pdf>

Worth mentioning is the fact that during 2008, 35 Gt of CO<sub>2</sub> were emitted from earth, (Kahn, 2009, Reuters 2009), without counting the CO<sub>2</sub> derived from the animal kingdom, while the increase of CO<sub>2</sub> concentration in the atmosphere was only 13 Gt. (1.66 ppmv, Mauna Loa Observatory). As a result only 37% of the emitted CO<sub>2</sub> from mother earth stays in the atmosphere. The remaining 22 Gt (63%) returns to earth. And this is done every year. According to NASA (Orbiting Carbon Observatories, OCO) some 66% of the emitted 35Gt can be attributed to hydrocarbons.

This implies that hydrocarbons contribute roughly 23Gt of CO<sub>2</sub> every year, roughly the amount that returns back to earth (22Gt), and hence we have more than one source contributing to the atmospheric CO<sub>2</sub> concentration since the total sum is 35 Gt. The latter could have been easily deduced from the work done by the European scientists from 1812 to 1961, Beck 2007. Moreover, it is more than obvious that at least 10Gt of CO<sub>2</sub> derived from hydrocarbons, (43.5%), returns back to earth every year. In reality we do not know the exact CO<sub>2</sub> sources, hence their percent contribution, nor we know where the 22Gt of CO<sub>2</sub> is disappearing, For this reason, NASA as well as Canada (Spears, 2009) sent, early this year, satellites to resolve these questions.



If we take into account that during the last 50 years, atmospheric CO<sub>2</sub> concentration has increased by 70 ppmv, that is by 0,007%, and that according to NASA OCO 2009 data, 66% of this amount is derived from hydrocarbons, that is 0,0046%<sup>3</sup>, then one should wonder if this negligible amount (change of atmospheric air composition in the third decimal point) has caused the climatic change. Is it possible an increase of 2 or even 3 mole, of CO<sub>2</sub> in 100000 moles of other atmospheric gases (currently 39 CO<sub>2</sub> moles in 100000 moles to 41 or 42 CO<sub>2</sub> moles in 100000 moles), to cause a climatic change?


It is therefore seen that there is no relation between atmospheric CO<sub>2</sub> concentrations and small temperature perturbations. But even if there were correlations the influence of CO<sub>2</sub> concentration on temperature is very weak (Lindzen 2006). Using a logarithmic relationship between the addition of CO<sub>2</sub> to the atmosphere and radiative heating, Lindzen estimated that the 100 ppm post industrial increase in CO<sub>2</sub> concentration (280 ppm pre-industrial to today's 380 ppm) has already caused about 75% of the anticipated I K (+0.37°C) warming, and finally an additional warming of few tenths of a degree occurs.

Sorokhtin et. al., (2007) tried to explain the so called "Green House Effect" using the adiabatic theory. This theory is based on the observation that in the troposphere the heat transfer is mainly carried out by convection and the temperature distribution is close to adiabatic. Their reasoning is that air masses expand and cool while rising and compress and heat while descending. As a result, even if the CO<sub>2</sub> concentration in the atmosphere is doubled, that is going from 350 ppm to 700 ppm, the temperature at sea level will increase by 0.01 °C. However, Rutledge (2007) as shown in Figure 22, calculated that of CO<sub>2</sub> concentrations derived from burning fossil fuels can not exceed 450 ppm. An excellent discussion on the non existing relation between CO<sub>2</sub> concentration in the atmosphere and temperature is presented by Florides and Christodoulides (2009).

---

<sup>3</sup> CO<sub>2</sub> contribution from solid fuels is 40% of the 0,0046% that is less than 0,002%. The carbon capture sequestration (ccs) aims at reducing this 0,002% to 0,001%.

## Atmospheric CO<sub>2</sub> Concentration with Various Carbon Emission Scenarios (2005 – 2400) Caltech, 2008



Copyright J. D. Hughes GSR Inc, 2008

(data from rutledge.caltech.edu, October, 2008)

**Figure 22.** Atmospheric CO<sub>2</sub> concentrations with various carbon emission scenarios from 2005 AD to 2400 AD, Caltech 2008. Data from rutledge.caltech.edu, October 2008. ( Copyright, Hughes, 2009) Worth noticing is the background of 325 ppmv atmospheric CO<sub>2</sub> concentration without the use of hydrocarbons, the same as Beck 2007 and the same if all Calendar's, 1985,data are taken into account

### 5. Relation of Temperature to Solar Activity

The question still remaining is how the small temperature perturbations can be explained when fluctuating between -0.5°C, “The Little Ice Age”, to + 0.35°C, today's increase. The answer can be found in the Sun's activity, the so called sunspots and solar winds (Svensmark-Friss-Christensen, 1997; Svensmark, 1998; 2007) (Figures 23 and 24).




Figure 23. An image of a large eruptive prominence emerging from the solar surface along with an artist illustration of the Sun-Earth connection, Marhavilas, 2008





Figure 24. Sun spot number variations from 1700 AD to 1995 AD. [www.ngdc.noaa.gov/stp/SOLAR/SSN/ssn.html](http://www.ngdc.noaa.gov/stp/SOLAR/SSN/ssn.html).

Sunspots are storms on the sun's surface that are marked by intense magnetic activity and host solar flares and hot gassy ejections from the sun's corona. The number of spots on the sun cycles over time, reaching a peak, the so-called Solar Maximum, every 11 years, Figure 25. Solar winds, according to NASA's Marshall Space Flight Center, consist of magnetized plasma flares and in some cases are linked to sunspots. They emanate from the sun and influence the amount of galactic dust (Murad and Williams, 2002; Landgraf, 2003) Figure 26, which may in turn affect atmospheric phenomena on Earth, such as cloud cover.




**Figure 25.** Eleven (11) year sunspot cycle from 1995 to present. The fluctuation of sunspot numbers is characteristic, Hathaway, 2008. The correlation with temperature drop is characteristic, Figure 20.



**Figure 26. Picture of galactic dust emanating from Galaxy M104, Hubble Heritage Gallery Images, 2009. The Distance of M104 from earth is 28 million light years, it contains 800 billion suns and its diameter is estimated at 50000 light years.**


It is calculated that when Solar activity is at a minimum, such as in 2009, 40000 tons of galactic dust/space debris (ESA/NASA mission Ulysses) reach the earth's atmosphere inducing the condensation of water vapours which leads to clouds formation (Svensmark et al., 2007). Clouds in turn affect the variations seen in temperatures (Shaviv and Weizer, 2003), Figure 27.



**Figure 27. Celestial drive of Phanerozoic Eon climate?. The influence of cosmic ray flux related to temperature change for the last 500 million years with a cycle of about 200 million years, Wilson's cycles, Shaviv and Weiser, 2003**

So the emphasis is on water vapour, which is the number one greenhouse gas, and not CO<sub>2</sub> whose concentration in the atmosphere is 100 times less than water vapour, H<sub>2</sub>O. The correlation between sunspot number, the solar activity proxies and the <sup>10</sup>Be isotope concentration, which is an indicator of the amount of the galactic dust reaching earth, from 1600 AD until 2000 AD, is presented in Figure 28 (Beer et. al., 1994; Hoyt and Schatten 1998a; 1998b).





**Figure 28. Solar proxy activities based upon <sup>10</sup>Be concentration found in Dye-3, Greenland ice core. Beer et. al., 1994. <sup>10</sup>Be originates from the incoming galactic dust. Relation to Sunspot number from 1600 AD to 2000 AD, Hoyt and Schatten 1998a, 1998b**

Another research work by McLean et al., 2009, shows that the surge in global temperature since 1977 can be attributed to a climate shift in the Pacific Ocean, that is the relationship between El Niño South Oscillation (ENSO) effect and global temperature. The available data strongly suggests that future global temperatures will continue to change in response to ENSO cycling, solar radiation and volcanic activities.

## **6. Other Climatic anomalies**

Various climatic changes such as the Dansgaard-Oeschger (DO) event have been recognized from ice cores taken from Greenland (GRIP/GISP2) which go back to the end of the last interglacial period, the Eemian Interglacial time. These events seem to have been globally synchronous, and they lasted 1500 years (Voelker, 2003). A lower frequency Bond cycle (Bond et al., 1999; Schulz, 2002; Braun et. al., 2005) is characterized by unusually cold conditions that take place during the cold DO phase, the subsequent Heinrich event and the rapid warming phase that follows each Heinrich event (Heinrich 1988; Bond et.al.,

1992; Grousset et. al., 2000). During each Heinrich event, massive fleets of icebergs are released into the North Atlantic, carrying rocks picked up by the glaciers far out to sea. Heinrich events are marked in marine sediments by conspicuous layers of iceberg-transported rock fragments.

Many of the transitions in the DO and Bond cycles were rapid and abrupt, and they are being studied intensively by paleoclimatologists and Earth system scientists to understand the driving mechanisms of such dramatic climatic variations that are not CO<sub>2</sub> driven. These cycles now appear to result from interactions between the atmosphere, oceans, ice sheets, and continental rivers that influence thermohaline circulation (the pattern of ocean currents driven by differences in water density, salinity and temperature rather than wind). Thermohaline circulation, in turn, controls ocean heat transport, such as the Gulf Stream which affects the climate of Northern Europe.

Climate, therefore, is driven by many natural processes which operate at many time scales with many scales of influence (Gerhard 2001), Figure 29. The mere fact that during the Huronian Glaciation, 2.4 billion years to 2.1 billion years, Paleoproterozoic Era and during the Cryogenian Period, 800 million years to 635 million years ago, Neoproterozoic Era, Earth was totally covered by snow, Tjeerd, 1994, Rieu et. al., 2007, Table 2, Figure 30, while atmospheric CO<sub>2</sub> concentrations were 10 times (Kan and Riding, 2007) to 200 times (Kaufman and Xiao, 2003) higher than today's, and the fact that during the Phanerozoic Era we have glaciations periods irrespective of atmospheric CO<sub>2</sub> concentrations, proves that there are many more natural drivers, besides the miniscule increases of atmospheric CO<sub>2</sub>, sunspots and solar winds and Milankovitch cycles, that influence Earth's climate.

**Table 2. Major glacial periods in earth's history  
Glaciations Periods During Paleoproterozoic and Neoproterozoic Ages  
and the Phanerozoic Eras.**

[www.en.Wikipedia.org/wiki/Timeline\\_of\\_glaciation](http://www.en.Wikipedia.org/wiki/Timeline_of_glaciation).

Name	Period (Ma)	Period	Era
<a href="#">Quaternary</a>	30 - present	<a href="#">Neogene</a>	<a href="#">Cenozoic</a>
<a href="#">Karoo</a>	360 - 260	<a href="#">Carboniferous</a> and <a href="#">Permian</a>	<a href="#">Paleozoic</a>
<a href="#">Andean-Saharan</a>	450 - 420	<a href="#">Ordovician</a> and <a href="#">Silurian</a>	<a href="#">Paleozoic</a>
<a href="#">Cryogenian</a> (or Sturtian-Varangian)	800 - 635	<a href="#">Cryogenian</a>	<a href="#">Neoproterozoic</a>
<a href="#">Huronian glaciation</a>	2100 - 2400	<a href="#">Siderian</a> and <a href="#">Rhyacian</a>	<a href="#">Paleoproterozoic</a>






Figure 29. Geologic constraints on global climate variability. Natural climate drivers ranked by intensity and duration. Human interventions meteorite impacts, volcanic eruptions, El Nino and others are considered the 4<sup>th</sup> Order affecting climatic variability, Gerhard, 2004




Figure 30. Major glacial periods in earth's history. [www.en.Wikipedia.org/wiki/Timeline\\_of\\_glaciation](http://www.en.Wikipedia.org/wiki/Timeline_of_glaciation).

## **7. Conclusions**

- I. Climatic changes have been the subject of detailed studies for the last 200 years, that is since the 1800s and not only in the last 20 years.**
- II. Climatic changes were the norm throughout the Phanerozoic Era.**
- III. At least 100 glacial and interglacial periods have been measured during the Quaternary Period using Geomagnetic Polarity data.**
- VI. Climatic Changes during the Quaternary Period were not caused by Homo sapiens but by natural climate drivers.**
- V. During the Holocene Epoch and prior to 1850 AD many climatic changes were identified: The Climatic Optimum from 8000 BC to 4000 BC, with temperatures 1°C to 2°C above the optimum Mean Annual Temperature of 15°C; The Roman Warm Period from 450 BC to 150 AD with temperatures over 2.5°C above the optimum Mean Annual Temperature; and the Medieval Warm Period from 900 AD to 1400 AD with temperatures over 0.4°C above the optimum Mean Annual Temperature. Homo sapiens who appeared 60000 years ago did not influence these climatic changes. The recent temperature increase of +0.35°C above the optimum Mean Annual temperature of 15°C from 1980 to 1998 can not be used to forecast catastrophic events, since from 1999 to date temperatures show a decreasing tendency. In addition, much higher temperatures have been recorded for longer time during the Holocene period without any damage to either earth or its population.**
- VI. The data, so far, do not support the relation between atmospheric CO<sub>2</sub> and temperature or other climatic changes. By looking into the more reliable and thus far overlooked chemical CO<sub>2</sub> methods for determining atmospheric concentrations one can not be positive about a relationship between temperature difference and CO<sub>2</sub> concentration. The adiabatic theory suggests that global warming, and hence climatic changes, due to atmospheric CO<sub>2</sub> concentrations is impossible.**
- VII. Climatic changes are very difficult to assess because there are many natural climate drivers with various intensities and different durations. So, earth's climate system can not be a function of one and only one factor, namely atmospheric CO<sub>2</sub> concentration.**

## 8. References

Agassiz, L. 1840. Etude sur les glaciers, Nauchatel. Digital book on Wikisource. [en.wikipedia.org/wiki/Louis\\_Agassiz](http://en.wikipedia.org/wiki/Louis_Agassiz)

Allen, J.R.M., Long, A.J., Ottley, C.J., Pearson, D.G., Huntley, B. 2007. Holocene climate variability in northernmost Europe. *Quaternary Science Reviews* 26: 1432-1453.

Arrhenius, S. 1896. "On the Influence of Carbonic Acid in the Air Upon the Temperature of the Ground." *Philosophical Magazine* 41: 237-76.

Arrhenius, S. 1901. "Über die Wärmeabsorption Durch Kohlensäure und Ihren Einfluss auf die Temperatur der Erdoberfläche." *Förhandlingar Svenska Vetenskapsakademiens* 58: 25-58.

Atmospheric CO<sub>2</sub> at Mauna Loa Observatory, 2009. Scripps Institute of Oceanography and US Department of Commerce, National Oceanic and Atmospheric Administration, Earth System Research Laboratory, Gas Monitoring Data (NOAA ESRL/GMD). [Http://en.wikipedia.org/wiki/File: CO<sub>2</sub>-Mauna-Loa .png](http://en.wikipedia.org/wiki/File:CO2-Mauna-Loa.png)

Barnola, J.-M., Raynaud, D., Korotkevich, Y. S., Lorius, C. 1987. Vostok ice core provides 160,000-year record of atmospheric CO<sub>2</sub>. *Nature* 329:408-14.

Barnola, J.-M., Raynaud, D., Neftel, A., Oeschger, H. 1983. Comparison of CO<sub>2</sub> measurements by two laboratories on air from bubbles in polar ice. *Nature* 303:410-13.

Barnola, J.-M., Pimienta, P., Raynaud, D., Korotkevich, Y.S. 1991. CO<sub>2</sub>-climate relationship as deduced from the Vostok ice core: A re-examination based on new measurements and on a re-evaluation of the air dating. *Tellus* 43(B):83- 90.

Barendregt, R.W., Duk-Rodkin, A. 2004 "Chronology and Extent of Late Cenozoic Ice Sheets in North America: A magnetostratigraphic Assessment" in *Quaternary Glaciations-Extent and Chronology, Part II*, editors, J. Ehlers and P.L. Gibbard. *Quaternary Science Reviews*. Elsevier, p. 1-7.

Beck, E. G. 2007. 180 years of atmospheric CO<sub>2</sub> gas analysis by chemical methods. *Energy and Envir.* v.18 No 2 pp. 259-282.

Beer, J., Baumgartner, St., Dittrich-Hannen, B., Hauenstein, J., Kubik, P., Lukaczyk, Ch., Mende, W., Stellmacher, R., Suter, M. 1994. Solar Variability Traced by Cosmogenic Isotopes in *The Sun as a Variable Star: Solar and Stellar Irradiance Variations* (eds. J.M. Pap, C. Fröhlich, H.S. Hudson and S.K. Solanki), Cambridge University Press, 291-300

Beerling, D.J., McElwain, J.C., Osborne, C. P. 1998. Stomatal responses of the 'living fossil' *Ginkgo biloba* L. to changes in atmospheric CO<sub>2</sub> concentrations. *Journal of Experimental Botany* 49: 1603-1607.

Berglund, B.E. 2003. Human impact and climate changes -- synchronous events and a causal link? *Quaternary International* 105: 7-12.

Berner R. A., Kothavala Z. GEOCARB III: 2002. A revised model of atmospheric CO<sub>2</sub> over Phanerozoic time. IGBP PAGES/World Data Center for Paleoclimatology, Data Contribution Series #2002-051. NOAA/NGDC Paleoclimatology Program, Boulder CO, USA; 2001.  
[http://www1.ncdc.noaa.gov/pub/data/paleo/climate\\_forcing/trace\\_gases/phanerozoic](http://www1.ncdc.noaa.gov/pub/data/paleo/climate_forcing/trace_gases/phanerozoic)

Blundell, A., Barber, K. 2005. A 2800-year palaeoclimatic record from Tore Hill Moss, Strathspey, Scotland: the need for a multi-proxy approach to peat-based climate reconstructions. *Quaternary Science Reviews* 24: 1261-1277.

Bond, G., Heinrich, H., Broecker, W., Labeyrie, L., Mcmanus, J., Andrews, J., Huon, S., Jantschik, R., Clasen, S., Simet, C. 1992. "[Evidence for massive discharges of icebergs into the North Atlantic ocean during the last glacial period](#)". *Nature* 360 (6401): 245–249.

Bond, G.C., Showers, W., Elliot, M., Evans, M., Lotti, R., Hajdas, I., Bonani, G., Johnson, S. 1999. "[The North Atlantic's 1–2 kyr climate rhythm: relation to Heinrich events, Dansgaard/Oeschger cycles and the little ice age](#)". in Clark, P.U., Webb, R.S., Keigwin, L.D.. *Mechanisms of Global Change at Millennial Time Scales*. Geophysical Monograph. American Geophysical Union, Washington DC. pp. 59–76. [ISBN 0-87590-033-X](#)

Braun, H., Christl, M., Rahmstorf, S., Ganopolski, A., Mangini, A., Kubatzki, C., Roth, K. and B. Kromer. 2005. Possible solar origin of the 1,470-year glacial climate cycle demonstrated in a coupled model. *Nature*. 438: 208-211.  
[doi:10.1038/nature04121](https://doi.org/10.1038/nature04121)

Briffa, K.R., Osborn, T.J., Schweingruber, F.H., Harris, I.C., Jones, P.D., Shiyatov, S.G., Vaganov, E.A. 1998. A low-frequency Temperature Variations from a Northern Tree Ring Density Network. *Journal of Geophysical Research*, 106 D3 pp. 2929-2941

Briffa K.R., Jones, P.D., Schweingruber, F.H., and Osborn, T. J. 1998b. "Influence of volcanic eruptions on northern hemisphere summer temperature over the past 600 years." *Nature*, Volume 393, pp. 450-455.

Broecker, W. S., 2001. Was the Medieval Warm Period global?, *Science*, 291, pp.1497-1499

Brown, Dale Mac Kenzie, 2000. The fate of Greenland's Viking. Time-Life's Book.[www.archaeology.org/online/features/Greenland/](http://www.archaeology.org/online/features/Greenland/)

Caillon N, Severinghaus J.P., Jouzel J., Barnola, J. M., Kang, J., Lipenkov, V.Y. 2003. Timing of atmospheric CO<sub>2</sub> and Antarctic temperature changes across termination III. *Science*, 299 (5613) pp. 1728–1731

Callendar, G.S., 1958. On the amount of carbon dioxide in the atmosphere *Tellus*, Vol. 10, pp. 243-248.

Cande, S. C., Kent, D. V. 1995. Revised calibration of the geomagnetic polarity timescale for the Late Cretaceous and Cenozoic. *Journal of Geophysical Research*, B, Solid Earth and Planets, v.100, pp 6093- 6095.

Comiso, J.C. 2000. Variability and trends in Antarctic surface temperatures from *in situ* and satellite infrared measurements. *Journal of Climate* 13: 1674-1696.

Croll, J. 1864. "On the Physical Cause of the Change of Climate During Geological Epochs." *Philosophical Magazine* 28: 121-37.

Czerney, Franz von. 1881. Die Veränderlichkeit des Klimas und Ihre Ursachen. Vienna

Dansgaard, W., Johnsen, S.J., Moller J. 1969. One thousand centuries of climatic record from the Camp Century on the Greenland Sheet. *Science*, v. 166 (3903), 377-381p

D' Arrigo, R., Mashig, E., Frank, D., Jacoby, G. and Wilson, R. 2004. Reconstructed warm season temperatures for Nome, Seward Peninsula, Alaska. *Geophysical Research Letters* 31: 10.1029/2004GL019756

Doran, P.T., Prisco, J.C., Lyons, W.B., Walsh, J.E., Fountain, A.G., McKnight, D.M., Moorhead, D.L., Virginia, R.A., Wall, D.H., Clow, G.D., Fritsen, C.H., McKay, C.P. and Parsons, A. N. 2002. Antarctic climate cooling and terrestrial ecosystem response. *Nature* advance online publication, 13 January 2002 (DOI 10.1038/nature710).

EPICA community members 2004. Eight glacial cycles from an Antarctic ice core, *Nature* 429:6992, 623-628, doi:10.1038/nature02599.

Falcon-Lang, H.J. 2005. Global climate analysis of growth rings in woods, and its implications for deep-time paleoclimate studies. *Paleobiology* 31(3):434-444. 2005

Fisher, H., Wahlen, M., Smith, J., Mastoianni, D., Deck, B. 1999. Ice core records of atmospheric CO<sub>2</sub> around the last three Glacial terminations. *Science*, v 283, 1712-1714p

Fleitmann, D., Burns, S.J., Neff, U., Mudelsee, M., Mangini, A., Matter, A. 2004. Palaeoclimatic interpretation of high-resolution oxygen isotope profiles derived from annually laminated speleothems from Southern Oman. *Quaternary Science Reviews* 23: 935-945.

Florides, G. A., Christodoulides, P. 2009. Global warming and carbon dioxide through sciences. *Environmental International*, v. 35, pp. 390-401

Fourier, J. 1824. "Remarques Générales sur les Températures Du Globe Terrestre et des Espaces Planétaires." *Annales de Chimie et de Physique* 27: 136-67. Translation by Ebeneser Burgess, "General Remarks on the Temperature of the Earth and Outer Space," *American Journal of Science* 32: 1-20 (1837).

Fourier, J. 1827. "Mémoire sur les Températures du Globe Terrestre et des Espaces Planétaires." *Mémoires de l'Académie Royale des Sciences* 7: 569-604

Franklin, B. 1784. "Meteorological Imaginations and Conjectures (Paper Read 1784)." *Memoirs of the Literary and Philosophical Society of Manchester* 2nd ed., 1789: 373-77. Reprinted *Weatherwise* 35, 262 (1982).

Friedli, H., Lotscher, H., Oeschger, H., Siegenthaler, U., Stauffer, B. 1986. "Ice core record of the  $^{13}\text{C}/^{12}\text{C}$  ratio of atmospheric  $\text{CO}_2$  in the past two centuries." *Nature*, Vol. 324, pp. 237-238.

Friss-Christiensen, E., Lassen, K. 1991. Length of the solar cycle: An indicator of solar activity closely associated with climate. *Science*, 254, pp.698-700

Galaxy M104. Hubble Heritage Gallery Images, 2009. www. Heritage.stsci.edu/..../galindex.html

Gerhard, L. C. 2001. Natural processes are the most significant climate drivers. [www.geocraft.com/VWFossils/.../Gerhard\\_Climate\\_Change.pdf](http://www.geocraft.com/VWFossils/.../Gerhard_Climate_Change.pdf)

Grousset, F.E., Pujol, C., Labeyrie, L., Auffret, G., Boelaert, A. (2000-02-01). "Were the North Atlantic Heinrich events triggered by the behaviour of the European ice sheets?" (abstract). *Geology* 28 (2): 123–126. doi:10.1130/0091-7613(2000)28<123:WTNAHE>2.0.CO;2. <http://geology.geoscienceworld.org/cgi/content/abstract/28/2/123>

Grudd, H., Briffa, K. R., Karlen, W., Bartholin, T.S., Jones, P. D. and Kromer, B. 2002. A 7400-year tree-ring chronology in northern Swedish Lapland: natural climatic variability expressed on annual to millennial timescales. *The Holocene* 12: 657-665.

Hadley Meteorological Center, 2008. HadCRUT 3. Global temperature record. Met Office Hadley Center for Climatic Prediction and Research. [www.metoffice.gov.uk/corporate/pressoffice/20081216.html](http://www.metoffice.gov.uk/corporate/pressoffice/20081216.html) html-27K

Hathaway, D. 2008. What is wrong with the Sun? (Nothing). [science.nasa.gov/.../11\\_Jul\\_solarcycleupdate.html](http://science.nasa.gov/.../11_Jul_solarcycleupdate.html)

Hayes, J. D., Imbrie, J., Shackleton, N. J. 1976. "Variations in the Earth's Orbit: Pacemaker of the Ice Ages." *Science*, 194: no 4270, pp1121-32.

Heikkila, M., Seppa, H. 2003. A 11,000-yr palaeotemperature reconstruction from the southern boreal zone in Finland. *Quaternary Science Reviews* 22: 541-554.

Heinrich, H. 1988. Origin and consequences of cyclic ice rafting in the Northeast Atlantic Ocean during the past 130,000 years, *Quaternary Research*, 29, 142-152

Holmgren, K., Karlen, W., Lauritzen, S.E., Lee-Thorp, J. A., Partridge, T. C., Piketh, S., Repinski, P., Stevenson, C., Svanered, O. and Tyson, P. D. 1999. A 3000-year high-resolution stalagmite-based record of paleoclimate for north-eastern South Africa. *The Holocene* 9: 295-309.

Holmgren, K., Tyson, P. D., Moberg, A., Svanered, O. 2001. A preliminary 3000-year regional temperature reconstruction for South Africa. *South African Journal of Science*, 99, pp. 49-51.

Hormes, A., Karlen, W., Possnert, G. 2004. Radiocarbon dating of palaeosol components in moraines in Lapland, northern Sweden. *Quaternary Science Reviews* 23: 2031-2043.

Hoyt, D. V., Schatten, K.H. 1998a. "Group sunspot numbers: A new solar activity reconstruction. Part 1.". *Solar Physics* 179: 189-219.

Hoyt, D. V., Schatten K.H. 1998b. "Group sunspot numbers: A new solar activity reconstruction. Part 2.". *Solar Physics* 181: 491-512.

Hughes, J. D. 2009. The energy sustainability dilemma: Powering the future in a finite world. Global Sustainability Research, Inc. Contact: dave-hughes@xplornet.com

Idso, C.D., Idso, K.E. 2000. *The Greening of the American West*. Center for the Study of Carbon Dioxide and Global Change, Tempe, AZ, USA.

Jaworowski, Z., Segalstad, T. V., Ono, N. 1992. Do glaciers tell a true atmospheric CO<sub>2</sub> story? *The Science of the Total Environment*, 114: p. 227-284.

Jones, P. 2008. Global temperature record. [www.cru.uea.ac.uk/cru/info/warming/](http://www.cru.uea.ac.uk/cru/info/warming/) and Brohan, P., J.J. Kennedy, I. Harris, S.F.B. Tett and P.D. Jones, 2006: Uncertainty estimates in regional and global observed temperature changes: a new dataset from 1850. *J. Geophysical Research* 111, D12106, [doi:10.1029/2005JD006548](https://doi.org/10.1029/2005JD006548)

Jiang, H., Seidenkrantz, M-S., Knudsen, K. L., Eiriksson, J. 2002. Late-Holocene summer sea-surface temperatures based on a diatom record from the north Icelandic shelf. *The Holocene* 12: 137-147.

Kah, L. C., Riding, R. 2007. Mesoproterozoic carbon dioxide levels inferred from calcified cyanobacteria. *Geology*, v.35. no 9, pp. 799-802

Kahn, M. 2009. Forests absorb 20% of fossil fuel emissions: [study.www.reuters.com/article/...../idUSTRE5IH5KE20092018](http://www.reuters.com/article/...../idUSTRE5IH5KE20092018)

Kaufman, A. J., Xiao, S. 2003. High CO<sub>2</sub> levels in the Proterozoic atmosphere estimated from analyses of individual microfossils. *Nature*, 425 (6955), pp. 279-282

Keigwin, L. D. 1996. The Little Ice Age and Medieval Warm Period in the Sargasso Sea. *Science*, 274: pp. 1504-1508.

Kelly, P. F., Kobes, R., Kunstatter, G. 1994. Parametrization invariance and the resolution of the unitary gauge puzzle *Phys. Rev. D* 50, pp.7592 – 7602

Khilyuk, K. Chillingar G. V., 2003. Global warming: are we confusing cause and effect. *Energy Sources* 25, pp. 357-370

Khilyuk, K. Chillingar G. V., 2006. Global forces of nature driving the Earth's climate. Are humans involved? *Environmental Geology* 50: pp. 899-910.

Krüger, T. 2008. Die Entdeckung der Eiszeiten. *Internationale Rezeption und*

Konsequenzen für das Verständnis der Klimageschichte, Basel 2008, [ISBN 978-3-7965-2439-4](#), pp.220-223, pp. 223-224, pp. 540-542

Köppen, W. 1873. "Über Mehrjährige Perioden der Witterung, Insbesondere Über die 11jährige Periode der Temperatur." *Zeitschrift der Osterreichischen Gesellschaft für Meteorologie* 8: 241-48, 141-50.

Kouwenberg, L.L.R., McElwain, J.C., Kurschner, W.M., Wagner, F., Beerling, D.J., Mayle, F.E. and Visscher, H. 2003. Stomatal frequency adjustment of four conifer species to historical changes in atmospheric CO<sub>2</sub>. *American Journal of Botany* 90: pp. 610-619

Kouwenberg, L. 2004. Application of Conifer Needles in the Reconstruction of Holocene CO<sub>2</sub> Levels. PhD Thesis, University of Utrecht; 2004.  
<http://www.bio.uu.nl/~palaeo/Personneel/Lenny/artikellinks/full.pdfN>.

Kouwenberg, L., Wagner, R., Kurschner, W., Visscher, H. 2005. Atmospheric CO<sub>2</sub> fluctuations during the last millennium reconstructed by stomatal frequency analysis of *Tsuga heterophylla* needles. *Geology* v.33 pp.33–36.

Kurschner, W.M., Burgh, van der, J., Visscher, H. and Dilcher, D.L., 1996. "Oak leaves as biosensors of late Neogene and early Pleistocene paleoatmospheric CO<sub>2</sub> concentrations." *Marine Micropaleontology*, Vol. 27, pp. 299-312.

Lamb, H., H. 1995. *Climate, History and the Modern World*. Publisher : Routledge , 464 p. ISBN: 13: 9780415127356

Landgraf, M. 2003. Galactic dust storm enters Solar System. [www. Mewscientist.com/.../dn4021-galactic-dust-storm-enters-solar-system.html](http://www.Mewscientist.com/.../dn4021-galactic-dust-storm-enters-solar-system.html)

Le Roy Ladurie, E. 1988. *Times of feast, times of famine: a history of climate since the year 1000*. Farrar-Straus and Giroux, Publisher, ISBN: 13: 9780374521220

Loehle, C. 2004. Climate change: detection and attribution of trends from long-term geologic data. *Ecological Modelling* 171: 433-450.

Loehle, C. 2005. Estimating climatic timeseries from multi-site data afflicted with dating error. *Mathematical Geology*, 37, pp. 127-140.

Loehle, C. 2007. A 2000 Year Global Temperature Reconstruction based on Non-Tree ring Proxy Data. *Energy & Environment* 18:1049-1058.

Loehle, C., and J.H. McCulloch. 2008. Correction to: A 2000-year global temperature reconstruction based on non-treering proxies. *Energy & Environment* 19(1): 93-100.

Linderholm, H.W., Gunnarson, B. E. 2005. Summer temperature variability in central Scandinavia during the last 3600 years. *Geografiska Annaler* 87A: 231-241.

Lindzen R. 2005. Understanding common climate claims. International Seminar on Nuclear War and Planetary Emergencies—34th Session, Erice, Italy, 19–24



August, 2005. Proceedings. World Federation of Scientists; 2006.  
<http://www.worldscibooks.com/environsci/6076.html>

Mann, M. E., Bradley, R. S., Hughes, M. K. 1998. Global-scale temperature patterns and climate forcing over the past six centuries. *Nature* 392: 779-787.

Mann, M. E., Bradley, R. S., Hughes, M. K. 1999. Northern hemisphere temperatures during the past millennium: inferences, uncertainties, and limitations. *Geophysical Research Letters*, 26, pp. 759-762.

Marhavilas, P. K. 2008. The space as a natural laboratory for electrotecnies. *Journal of Engineering Science and Technology*, v 1, pp.9-18.

Mariolakos, I. D. 2008. Water and Environment. A Geo-Mythological Approach. AQUA 2008. 4th Intl. Exhibition of Water and Environment.  
[www.mio-ecsde.org/filemgmt/visit.php2?lid=364](http://www.mio-ecsde.org/filemgmt/visit.php2?lid=364)

Milankovitch, M. 1930. "Mathematische Klimalehre und Astronomische Theorie der Kilimaschwankungen." In *Handbuch der Klimatologie*, edited by W. Köppen and R. Geiger, Vol. 1, Pt. A, pp. 1-176. Berlin: Borntraeger

Moberg, A., D.M. Sonechkin, K. Holmgren, N.M. Datsenko, and W. Karlén, 2005. Highly variable Northern Hemisphere temperatures reconstructed from low- and high resolution proxy data. *Nature*, 433, pp. 613-617.

Munroe, J. S. 2003. Estimates of Little Ice Age climate inferred through historical rephotography, Northern Unita Mountains, U.S.A. *Arctic, Antarctic, and Alpine Research* 35: 489-498.

Murad, E., Williams, I. P. 2002. *Meteors in the Earth's Atmosphere: Meteoroids and Cosmic Dust and their Interaction with Earth's Upper Atmosphere*. Cambridge Univ. Press 322p. ISBN-13: 9780521804318, ISBN: 10: 0521804310

NASA's OCO satellite. Aims to Solve a Climate Change. *Scientific American* 2009 [www.scientificamerican.com/article.cfm?id=nasas...satellite](http://www.scientificamerican.com/article.cfm?id=nasas...satellite)

Neftel, A., Moor, E., Oeschger, H., Stauffer, B., 1985. "Evidence from polar ice cores for the increase in atmospheric CO<sub>2</sub> in the past two centuries." *Nature*, Vol. 315, pp. 45-47.

Olafsdottir, R., Schlyter, P., Haraldsson, H.V. 2001. Simulating Icelandic vegetation cover during the Holocene: Implications for long-term land degradation. *Geografiska Annaler* 83A: 203-215.

Petit, J.R., Basile, I., Leruyet, A., Raynaud, D., Lorius, C., Jouzel, J., M. Stievenard, M., Lipenkov, V.Y., Barkov, N.I., Kudryashov, B.B., Davis, M., Saltzman, E., Kotlyakov, V. 1997. Four climate cycles in Vostok ice core. *Nature* 387: 359-360

Petit, J.R., Jouzel, J., Raynaud, D., Barkov, N.I., Barnola, J-M., Basile, I., Benders, M., Chappellaz, J., Davis, M., Delayque, G., Delmotte, M., Kotlyakov, V.M., Legrand, M., Lipenkov, V.Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E.,

Stievenard, M. 1999. Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica. *Nature* 399: 429-436.

Pidwirny, M. 2006. "Earth's Climatic History". *Fundamentals of Physical Geography, 2nd Edition*. Date Viewed.  
<http://www.physicalgeography.net/fundamentals/7x.html>

Pisaric, M., Carey, S., Kokelj, S., Youngblut, D. 2007. Anomalous 20th century tree growth, Mackenzie Delta, Northwest Territories, Canada. *Geophys Res Lett* 34. doi:[10.1029/2006GL029139](https://doi.org/10.1029/2006GL029139)

Raymo, M. E. 1992. Global climate change: a three million year perspective. In: Kukla, G. and Went, E. (eds.), *Start of a Glacial, Proceedings of the Mallorca NATO ARW, NATO ASI Series I, Vol. 3, Springer-Verlag, Heidelberg*, p. 207-223.

Reuters, 2009. Orbital Set to Launch Orbiting Carbon Observatory Earth Science Satellite Aboard Taurus Rocket for NASA, Monday February 23, 2009

Rieu, R., Allen, P.A., Plotze, M., Pettke, T. 2007. Climatic cycles during a Neoproterozoic "snowball" glacial epoch" *Geology* v. 35(4): pp. 299–302

Royer D. L., Berner, R. A., Montañez, I. P., Tabor, N. J., Beerling, D. J. 2004. CO<sub>2</sub> as a primary driver of Phanerozoic climate. *GSA Today* 2004; 14(3). doi:10.1130/1052-5173.<<http://www.soest.hawaii.edu/GG/FACULTY/POPP/Royer%20et%20al.%202004%20GSA%20Today.pdf>>.

Ruddiman, W.F., Raymo, M. E., Martinson, D. G., Clement, B. M., Backman, J. 1989, Mid-Pleistocene evolution of Northern Hemisphere climate. *Paleoceanography*, v. 4, p. 353-412.

Rutledge, D., 2007. Hubbert's Peak, the coal question and climate change. [http://www.aspousa.org/proceedings/Houston/presentations/talk 20% for 20% ASPO % from Dave%20 Rutledge pdf](http://www.aspousa.org/proceedings/Houston/presentations/talk%20for%20ASPO%20from%20Dave%20Rutledge.pdf)

Schönwiese, C. 1995, "Klimaänderungen: Daten, Analysen, Prognosen", Springer, Heidelberg (Link to Amazon <http://www.amazon.com/Klima%C3%A4nderungen-Daten-Analysen-Prognosen-German/dp/354059096X>).

Schmidt, G. 2006. "[Medieval warmth and English wine](http://www.realclimate.org/index.php/archives/2006/07/medieval-warmth-and-english-wine/)". *RealClimate*. <http://www.realclimate.org/index.php/archives/2006/07/medieval-warmth-and-english-wine/>. Retrieved 2006-07-12.

Schneider, D.P., Steig, E.J., van Ommen, T.D., Dixon, D.A., Mayewski, P.A., Jones, J.M. and Bitz, C.M. 2006. Antarctic temperatures over the past two centuries from ice cores. *Geophysical Research Letters* 33: [10.1029/2006GL027057](https://doi.org/10.1029/2006GL027057)

Schulz, M. 2002. "On the 1470-year pacing of Dansgaard-Oeschger warm events". *Paleoceanography* 17 (2): 1014. doi:[10.1029/2000PA000571](https://doi.org/10.1029/2000PA000571)

Scotese, C. R. 2002. PALEOMAP Project, Arlington, Texas; 2002. <http://www.scotese.comN>.

Seppa, H., Birks, H., J., B. 2001. July mean temperature and annual precipitation trends during the Holocene in the Fennoscandian tree-line area: pollen-based climate reconstruction. *The Holocene* 11: 527-539

Seppa, H., Birks, H., J., B. 2002. Holocene climate reconstruction from the Fennoscandian tree-line area based on pollen data from Toskaljavri. *Quaternary Research* v.57, Issue 2, pp. 191-199

Shaviv, N.J., Weizer, J. 2003. Celestial driver of Phanerozoic climate? *GSA Today*, v.13, Issue 7, pp. 4-10

Siegenthaler, Urs., Stokes, T. F., Monnin, E., Luthi, D., Schwander, J., Stauffer, B., Raynaud, D., Barnola, J.-M., Fischer, H., Masson-Delmotte, V., Jouzel, J., 2005. Stable carbon cycle-climate relationship during the Late Pleistocene. *Science*, v 310, 1313-1317 p

Singer, F. S., Idso, G. 2009. Climate change reconsidered. The Heartland Institute, 880 p. ISBN-10: 1934791288

Slocum, G. 1955. Has the amount of carbon dioxide in the atmosphere changed significantly since the beginning of the twentieth century? *Month. Weather Rev.*, 1955(October): p. 225-231.

Soon, W., Baliunas, S. 2003. "Proxy Climatic and Environmental Changes of the Past 1000 Years." *Climate Research* 23: 89-110.

Sorokhtin, O. G., Chilingar, G.V., Khilyuk, L. F. 2007. Global warming and global cooling. Evolution of climate on earth. *Developments in Earth & Environmental Sciences*, Elsevier, 978-0-444-53815-5

Spears, T. 2009. Canadian mini-satellite may solve carbon puzzle. [www2.canada.com/calgaryherald/news/story.html?id](http://www2.canada.com/calgaryherald/news/story.html?id).

Spencer, R. 2007. 2000 years of Global Temperatures: The 2007-2008 Global cooling event. [www.drroy.spencer.com/](http://www.drroy.spencer.com/)-

Svensmark, H., Friss-Christensen, E. 1997. Variation of cosmic ray flux and global cloud relationships. *J. Atm. Solar-Terrest. Phys.* 59: 1225-1232.

Svensmark, H. 1998. "[Influence of Cosmic Rays on Earth's Climate](#)". *Physical Review Letters* 81: 5027–5030.

Svensmark, H, Pedersen, J. O. P., Marsh, N.D., Enghoff, M. B., Uggerhøj, U.I. 2007. "[Experimental evidence for the role of ions in particle nucleation under atmospheric conditions](#)". *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 463 (2078): 385–396.

Svensmark, H. 2007. "[Astronomy & Geophysics Cosmoclimate: a new theory emerges](#)". *Astronomy & Geophysics* 48 (1): 1.18–1.24.

Thompson, D.W.J. and Solomon, S. 2002. Interpretation of recent Southern Hemisphere climate change. *Science* 296: 895-899.

Tjeerd, H. van A. 1994. *New Views on an Old Planet: A History of Global Change* 2nd ed. Cambridge University Press, Cambridge, UK, p. 457 10: [ISBN 0521447550](#)

Tyndall, J. 1861. "On the Absorption and Radiation of Heat by Gases and Vapours." *Philosophical Magazine* ser. 4, 22: 169-94, 273-85.

Tyndall, J. 1863. "On Radiation through the Earth's Atmosphere." *Philosophical Magazine*, ser. 4, 25: 200-206.

Viau, A. E., Gajewski, K., Sawada, M. C., Fines, P. 2006. Millennial-scale temperature variations in North America during the Holocene. *Journal of Geophysical Research*, 111, D09102, doi: 10.1029/2005JD006031. (<http://www.lpc.uottawa.ca/data/reconstructions/index.html>)

Voelker, A., H., L. (2002). "Global distribution of centennial-scale records for Marine Isotope Stage (MIS) 3: a database". *Quaternary Science Reviews* 21: 1185–1212. doi:[10.1016/S0277-3791\(01\)00139-1](https://doi.org/10.1016/S0277-3791(01)00139-1)

Wagner, F., Sjoerd J. P., Bohncke, S. J. P., Dilcher, D.L., Kürschner, W.M., Geel, van B., Visscher, H. 1999. Century-Scale Shifts in Early Holocene Atmospheric CO<sub>2</sub> Concentration *Science*, v. 284. no. 5422, pp. 1971 – 1973

Wagner, F., Lenny L. R. Kouwenberg, L. L, R., van Hoof, T, B., Visscher, H. 2004. Reproducibility of Holocene atmospheric CO<sub>2</sub> records based on stomatal frequency. *Quaternary Science Review*, v.23, Issue 18-19, pp. 1947-1954

Wiki Answers, 2009. How reliable are air bubbles in ice core samples for determining historic levels of Carbon Dioxide in the atmosphere? [wiki.answers.com/.../How\\_reliable\\_are\\_air\\_bubbles\\_in\\_ice\\_core\\_samples\\_for\\_determining\\_historic\\_levels\\_of\\_Carbon\\_Dioxide\\_in\\_the...](http://wiki.answers.com/.../How_reliable_are_air_bubbles_in_ice_core_samples_for_determining_historic_levels_of_Carbon_Dioxide_in_the...)