

T H E E R A T O S T H E N E S E X P E R I M E N T

On the 20th of March 2019 Spring Equinox

On Wednesday, March 20th 2019, more than 20.000 students and 1500 teachers at more of 650 schools in Greece, just about the time the sun was at its highest point in the sky, using simple instrumentation, followed the steps and thoughts of the great scientist Eratosthenes, as happened in 240 BC. From northern Greece to Cyprus and from Imvros to Corfu, at noon on that day, the Eratosthenes' disciples and friends of learning, managed to calculate the circumference of the Earth, in an amazing fashion and with minimal error. Our students put sticks or other objects on the ground, measured their height and shadow's length and with simple mathematical reasoning and the aid of modern technologies (Google Earth, Sun Calc, etc.) calculated the circumference and the radius of the Earth.

The 2nd Junior High School of Amaliada, for fourth consecutive year, participated in this action with a group of students of the third grade. These students, members of the Erasmus+ group of our school, exchanged their data with the students of the collaborating I.I.S.S. Einaudi-Casaregis-Galileo Galilei school in Genoa, Italy, and calculated the Earth's radius with minimal error.

On that particular day, weather conditions were poor in the Amaliada region, with a lot of clouds obscuring the sun, making the whole activity challenging. Nevertheless, our science team were able to catch the necessary sun rays and proceed with the experiment.


Being part of the program European Schools Go Green, our students showed that the Sun and the light it emits can not only be used as an energy source but also as a tool for the evaluation of impossible things, like the Earth's radius.

Coincidentally, the measurement of the Earth's circumference by Eratosthenes links beautifully with Genoa's I.I.S.S. Einaudi-Casaregis-Galileo Galilei school, iconic symbol, Christopher Columbus, on his attempt to reach India from Europe by sail. Eratosthenes' calculations of the size of the Earth were very precise and very accepted during Columbus' time. Still, Columbus was convinced that the world was significantly smaller and that getting to Asia from Europe was feasible. Had he not rejected Eratosthenes' calculations, he would probably never have struggled his voyage, thus leaving the Americas undiscovered and changing the course of history.

Grigoris Vassilopoulos

WHO Eratosthenes WAS

Eratosthenes was a talented mathematician and geographer as well as an astronomer. He made several other important contributions to science. Eratosthenes devised a system of latitude and longitude, and a calendar that included leap years. He invented the armillary sphere, a mechanical device used by early astronomers to demonstrate and predict the apparent motions of the stars in the sky. He also compiled a star catalog that included 675 stars. His measurement of the circumference of Earth was highly respected in his day, and set the standard for many years thereafter. He may have also measured the distances from Earth to both the Moon and to the Sun, but the historical accounts of both deeds are, unfortunately, rather cryptic. A crater on Earth's Moon is named after Eratosthenes.



HOW HE DID IT

In 240 B.C., the Greek astronomer Eratosthenes made the first good measurement of the size of Earth. By noting the angles of shadows in two cities on the Summer Solstice, and by performing the right calculations using his knowledge of geometry and the distance between the cities, Eratosthenes was able to make a remarkably accurate calculation of the circumference of Earth. Eratosthenes lived in the city of **Alexandria**, near the mouth of the Nile River by the Mediterranean coast, in northern Egypt. He knew that on a certain day each year, the Summer Solstice, in the town of **Syene** (currently **Aswan**) in southern Egypt, there was no shadow at the bottom of a well. He realized that this meant the Sun was directly overhead in Syene at noon on that day each year. Eratosthenes knew that the Sun was never directly overhead, even on the Summer Solstice, in his home city of Alexandria, which is further north than Syene. He realized that he could determine how far away from directly overhead the Sun was in Alexandria by measuring the angle formed by a shadow from a vertical object. He measured the length of the shadow of a tall tower in Alexandria, and used simple geometry to calculate the angle between the shadow and the vertical tower. This angle turned out to be about 7.2 degrees. Next, Eratosthenes used a bit more geometry to reason that the shadow's angle would be the same as the angle between Alexandria and Syene as measured from the Earth's center. Conveniently, 7.2 degrees is 1/50th of a full circle ($50 \times 7.2^\circ = 360^\circ$). Eratosthenes un-

derstood that if he could determine the distance between Alexandria and Syene, he would merely have to multiply that distance by 50 to find the circumference of Earth! Eratosthenes had the distance between the two cities measured. His records show that the distance was found to be 5,000 stadia (approx. 800 Km). The stadion or stade (plural = stadia) was a common distance unit of the time. Unfortunately, there was not a universal, standard length for the stadion; so we don't know exactly which version of the stadion Eratosthenes used, and therefore are not exactly sure how accurate his solution was. He

may have been correct to within less than 1% (if used the Greek stadion that was approx. 155 meters), a remarkable accomplishment! Or, if it was actually a different stadion (if used the Italian attic stadion that was approx. 185 meters) that he used, he may have been off by about 16%.


The actual polar circumference of Earth is just a bit over 40,000 km.



Instructions

for

The measurement of the Radius of the Earth



Prerequisites

When two or more schools are collaborating, the experiment can be done on any day around the year, weather conditions permitting. In our case, it will be implemented on the same day for all schools, during the week of 18 – 22 March 2019. Let me remind you that the spring equinox is on the 21st of March.

All three schools should commence at local noon, where the sun is at its highest point in the sky. You can find out what time this is, either using suncalc.net or [stellarium](http://stellarium.stellarium.org) from stellarium.org, for the specific day of the action. Another idea is to find the time at which any object's shadow is minimum during the day. That would be local noon.


Because of its easternmost position, the 2nd Junior High School of Amaliada must perform the experiment first, whereas the Goethe High School of Kassel and Galileo Galilei Gymnasium in Genova one hour later, as the two schools lie roughly on the same meridian (*Fig.1 Eratosthenes_Maps.pdf*). At that time, the rotating Earth will carry the two schools at the same meridian as Amaliada was an hour ago.

The experiment

1. At local noon, a group of students should place a stick of known height h in a vertical position on the ground.
2. Measure the length of its shadow l .

ERATOSTHENES 2019

3. Calculate the angle theta θ , using the formula $\theta = \arctan \frac{l}{h}$ (see figure above).
4. Communicate with any of the two other schools, to obtain their relative angle θ .
5. Calculate the radial difference $\varphi = |\theta_1 + \alpha - (\theta_2 + \alpha)| = |\theta_1 - \theta_2|$.
6. Using **Google Earth** find the North – South distance **AB** between your school and the school whose angle θ you obtained earlier. The distances are given for comparison at *Fig.3 Eratosthenes_Maps.pdf*. By all means, you should let your students find themselves this data.
7. Calculate the circumference C of the Earth using the formula $C = \frac{AB \cdot 360^\circ}{\varphi}$
8. Find the radius of the Earth R_{\oplus} using $R_{\oplus} = \frac{C}{2\pi}$
9. Enjoy!!!



TEACHER'S GUIDE: THE ERATOSTHENES PROJECT

Overview

This activity is part of the World Year of Physics 2005, the centennial celebration of Einstein's "miracle year." In 1905, Einstein created three groundbreaking theories that provided the framework for much of the exciting physics of the last hundred years.

This activity enables students to measure the radius of Earth. Groups of students at two distant schools will take data and then collaborate, in essentially the same way that Eratosthenes measured Earth's circumference millennia ago in Egypt. A World Year of Physics website will enable each school to find a partner.

Objectives	2
Logistics	2
National Standards Addressed	2
Materials	2
Student's Guide	3
Additional Activities	6
Notes on the Activities	7
References	8

TEACHER'S GUIDE: THE ERATOSTHENES PROJECT

Objectives

- Describe the geometry of how sunlight strikes Earth at different latitudes
- Describe how the circumference of Earth was first measured millennia ago
- Describe how to determine local noon
- Measure the angle of the sun at local noon
- Collaborate with another school some distance away to determine the radius of Earth.

Logistics

Your class will need the measurements made at some other school located a substantial distance away, either north or south. You will enter your school's latitude and longitude on the Eratosthenes Project website, <<http://www.physics2005.org/events/eratosthenes/>>, and you will then be paired with another school. After the collaborating schools have shared measurements, and students have computed the radius of Earth, they will enter their results on the site. All results will be displayed on the site, and a grand average will be made of all the measurements to obtain the final value of Earth's radius determined by this project.

National Standards Addressed

This activity addresses the following Benchmarks and National Science Education Standards (NSES):

Benchmarks, K-12: **The Mathematical World**

Symbolic Relationships

When a relationship is represented in symbols, numbers can be substituted for all but one of the symbols and the possible values of the remaining symbol computed. . . .

Shapes

Distances and angles that are inconvenient to measure directly can be found from measurable distances and angles using scale drawings or formulas.

Models

The basic idea of mathematical modeling is to find a mathematical relationship that behaves in the same way as the objects or processes under investigation. A mathematical model may give insight about how something really works or may fit observations very well without any intuitive meaning.

Benchmarks, K-12: The Nature of Science

The Scientific Enterprise

The early Egyptian, Greek, Chinese, Hindu, and Arabic cultures are responsible for many scientific and mathematical ideas and technological inventions.

NSES, K-12: History and Nature of Science

Science as a Human Endeavor

Individuals and teams have contributed and will continue to contribute to the scientific enterprise.

Historical Perspectives

In history, diverse cultures have contributed scientific knowledge and technologic inventions.

NSES, K-12: Unifying Concepts and Processes

Developing Student Understanding

- Evidence, models, and explanation
- Constancy, change, and measurement

NSES, K-12: Science as Inquiry

Abilities necessary to do scientific inquiry

- Design and conduct scientific investigations
- Use technology and mathematics to improve investigations and communications
- Formulate and revise scientific explanations and models using logic and evidence

Materials

For each student

A copy of the *Student Activity Guide* (pages 3 to 5 of this document)

If students are doing the extension activities, see page 6.

For each group

- stick or dowel, about 60 centimeters (cm) long
- stiff cardboard (to provide a smooth, flat surface)
- meter stick
- pieces of blank paper
- pencil


To be shared by several groups

- carpenter's level
- tape

STUDENT'S GUIDE: THE ERATOSTHENES PROJECT

This activity is part of the World Year of Physics 2005, the centennial celebration of Einstein's "miracle year." In 1905, Einstein created three groundbreaking theories that provided the framework for much of the exciting physics of the last hundred years.

In this activity, you will work together with students at another school to measure the radius of Earth. You will use the same methods and principles that Eratosthenes used more than two thousand years ago.



Eratosthenes was a Greek living in Alexandria, Egypt, in the third century, BC. He knew that on a certain day at noon in Syene, a town a considerable distance to the south, the sun shone straight down a deep well. This observation meant that the sun was then directly overhead in Syene, as shown in Figure 1. Eratosthenes also knew that when the sun was directly overhead in Syene, it was *not* directly overhead in Alexandria, as shown in Figure 2. Notice that in both drawings, the sun's rays are shown as parallel.




Figure 1:
Light rays shining straight down a well in Syene at noon, when the sun is directly overhead. No shadow is cast.




Figure 2:
Sunlight shining down a well in Alexandria at noon, on the same day as the observation shown in Figure 1. The sun is not directly overhead. The gray bar at the bottom left shows the shadow cast by the side of the well. The angle of the sun's rays and the size of the shadow are exaggerated.

In Figure 2, the side of the well casts a shadow on the bottom. Eratosthenes used a shadow like this to find the circumference of Earth. When the sun was directly overhead in Syene, he measured the shadow of an object in Alexandria at noon. From the length of the shadow, the height of the object, and the distance between Syene and Alexandria, he calculated the circumference of Earth. His value agreed quite well with the modern one.

How Eratosthenes Found the Circumference of Earth




Figure 3:
The sun is directly overhead at noon at Syene (S). Alexandria is at point "A."

How did he do it, more than two thousand years ago? Take a look at Figure 3. Syene is represented by point "S," and Alexandria by point "A." In Figure 3, the arc length between S and A is d , and the angle corresponding to the arc SA is θ . The radius of Earth is R . As suggested above, let's assume that the sun's rays are parallel. Since the ray that strikes Syene, at point S, is perpendicular to the surface of Earth, the sun is directly overhead there.

When the sun was directly overhead in Syene, Eratosthenes measured the shadow of a tower in Alexandria at noon,¹ shown in Figure 4. Since both the tower at A, which is perpendicular to Earth's surface, and the ray of sunlight at point S both point to the center of Earth, and the rays of sunlight are parallel, the angle between the sunlight and the tower is equal to θ . (Alternate interior angles are equal.)




Figure 4:
The geometry of Eratosthenes' measurement. He measured the length of the tower and its shadow at noon at Alexandria. Then he determined the angle of sunlight with the vertical, which is the same as the angle subtended by Syene (S) and Alexandria (A) at the center of Earth.

The tower and its shadow form two sides of a right triangle, as shown in Figure 4. Although trigonometry hadn't yet been invented, Eratosthenes' procedure can be expressed in the language of trig as follows: The length of the shadow, the height of the tower (which he knew), and the angle θ , given here in degrees, are related by

$$\tan \theta = \frac{\text{length of shadow}}{\text{height of tower}} \quad (1)$$

Inverting $\tan \theta$ gives the value of θ . Using ratio and proportion, the arc length d is the same fraction of Earth's circumference C as θ is of 360 degrees.

$$\frac{d}{C} = \frac{\theta}{360} \quad (2)$$

Rearranging for the circumference C ,

$$C = (360/\theta)d \quad (3)$$

¹Some accounts say that Eratosthenes made measurements with a needle that cast a shadow onto the graduated inner surface of a hemispherical bowl. To relate as closely as possible to the method of this project, we will describe the experiment in terms of the shadow of a tower.

How You Can Find the Radius of Earth

Rather than find Earth's circumference, we suggest you find its radius, so that you can more easily compare your measured value with the accepted value.



Eratosthenes was lucky, because he knew of a place where the sun was directly overhead at noon on a certain day. Can you do the experiment even without that information? Fortunately you can, as shown in Figures 5 and 6. You will make the shadow measurements at your school and share results with a class from a school at another location. You can then do a subtraction to find the angle you need. Be sure to plan ahead—ideally, both schools make the measurement on the same day.




Figure 5: The geometry for measuring the radius of Earth using the data of two collaborating schools separated by a north-south distance d . Each will measure the angle of the sun, θ_A at one location and θ_B at the other, at local noon.

We need two points, A and B , separated by a north-south distance d , shown on Figure 5. The experiment will work best if d is as large as possible.

Take a look at Figure 6. Your school and the collaborating school are represented by the points A and B , and the angles θ_A and θ_B correspond to points A and B .




Figure 6: The relationship among the direction of sunlight, the sticks, and the two angles θ_A and θ_B .

At point A ,

$$\tan \theta_A = \frac{\text{length of shadow}}{\text{height of stick}} \quad (4)$$

and likewise at point B .

Figures 5 and 6 show that the angle corresponding to the arc AB is just the difference $\theta_B - \theta_A$. We can find the radius of Earth in the same way that Eratosthenes found the circumference in equation (3).

$$\frac{d}{2\pi R} = \frac{\theta_B - \theta_A}{360} \quad (5)$$

Rearranging and simplifying,

$$R = \frac{180d}{\pi(\theta_B - \theta_A)} \quad (6)$$

Making the Measurement at Local Noon

On any day, local noon is the instant when the sun reaches its highest point in the sky. To determine it, plant the stick in the ground, making sure the stick is vertical using a plumb bob or a carpenter's level. In the late morning, measure the shadow's length at regular time intervals. The shadow will get shorter as noon approaches, and then get longer again once noon has passed. The shortest length is what you will substitute into equation (4) above to find the value of θ_B or θ_A for your location.

ADDITIONAL ACTIVITIES

Below are three additional activities:

- How Shadows Change During the Day
- Shadows on Earth
- Latitude

How Shadows Change During the Day

If your students have not thought much about shadows, they might benefit by starting with this preliminary activity. Give each group a 5-cm straw piece, a sheet of 8½" x 11" paper, and some tape. Ask them to tape the straw so it stands in the center of the paper, as shown in Figure 7, and also to indicate the direction of north on one of the long sides. If you provide them with a compass, they can orient the paper.




Figure 7:
Five-centimeter straw mounted vertically on a piece of paper. Students predict and then measure the shadow of this straw at different times.

Their challenge is to imagine that this paper is set on level ground in sunlight and to predict the location and length of the shadow of the stick on the hour during the day. Discuss these predictions to bring out their thinking. Then have them do the experiment and compare their predictions and results.

Shadows on Earth

Materials for each group: five 4-cm straw pieces, tape, and a piece of 8½" x 11" paper.

Explain to students that they will make a model of shadows at different points on Earth. Have them draw a straight line across a piece of 8½" x 11" paper and tape the five straw pieces, equally spaced, along this line, so the straws stand straight up. Ask how the paper and straws could be a model of sticks placed at different locations on Earth (curve the paper, with the straws on the convex side).

Explain that to avoid damaging their eyes, they should never look directly at the sun. In sunlight, ask students how the paper and straws can model the Eratosthenes experiment. (Facing the sun, hold the paper at the ends of the long sides and curve it so the straws point out. Turn the paper to make the shadow of one straw disappear. Make this

straw point directly at the sun. The straw without a shadow models the well at Syene.) Have students describe what happens to the shadows of the other straws and relate the shadow of each one to its position. Ask students to relate these shadows to the shadows Eratosthenes used to measure Earth's circumference. See Figure 8.



Figure 8:
Model of shadows of sticks at noon, at different latitudes and the same longitude.

Extension Activity: Latitude

Once experimentation is complete and the results reported, you can have students relate the measurements they have made to the definition of latitude.

- Ask them to define latitude (the length of arc, or angle from the center of Earth, measured north or south from the equator).
- Referring them to Figure 3, ask them to assume that point *S* is on the equator. Ask on what day the sun would be directly overhead at noon at *S*. (In 2005, March 20, the vernal equinox, and September 22, the autumnal equinox; at the equinoxes; day and night have equal length, and the axis of Earth's spin is perpendicular to the line from Earth to the sun.)
- If your students made their shadow measurement on the vernal or autumnal equinox, the resulting angle would be equal to the latitude, as shown in Figure 3 (remember point *S* is on the equator). If possible, have them try to do this by measuring the shadow of a stick on or near March 20 or September 22.
- In the Eratosthenes experiment, the angle $(\theta_B - \theta_A)$ is the same as the difference in latitude of the two schools, so students could determine this difference immediately by subtracting the two latitudes of the collaborating schools. Of course, we want students to make measurements and compare, rather than look up the answer in an atlas. If students point this out, you can remind them that they are reenacting an historical experiment.

NOTES ON THE ACTIVITIES

Notes on Introduction

In the discussion of Figures 1 and 2, which show sunlight in wells at two different locations, we assume the following:

- The rays of sunlight are parallel (see next section for more detail)
- The sides of the well are vertical.

Notes on How Eratosthenes Found the Circumference of Earth

Here are Eratosthenes' assumptions:

Earth is a sphere. In fact, Earth bulges by about .3% at the equator, but we can safely neglect this difference.

The sun is very far away, so sunlight can be represented by parallel rays. The sun is indeed very far away, but it is not a point source, since its diameter is about 1/100 of the Earth-sun distance.




Figure 9:
Notice the penumbra—the partially illuminated region—at the end of the stick's shadow (drawing not to scale).

As shown in Figure 9, there is a penumbral region at the end of the shadow, a region only partially illuminated by the sun. If the stick is 1 meter (m) long, the penumbral region will be more than 1 centimeter (cm) wide, which limits the accuracy of the measurement of the shadow length. The penumbra size scales up with the length of the stick, so using a longer stick does not increase the accuracy of the measurement.

Alexandria is directly north of Syene: This is only approximately true. Find an atlas and compare the location of Alexandria and Aswan (built on the site of Syene).

You might ask students to comment on Eratosthenes' assumptions, considering that he was working more than 2,000 years ago.

Notes on How You Can Find the Radius of Earth

Ask students to discuss the distance d between the two schools. Is it better for d to be large or small? (large) Why? (The larger d , the larger the value of the angle $\theta_B - \theta_A$. The larger this angle, the smaller the *percentage* error in its measurement.)


Once you have obtained the name and location of the school you'll collaborate with, show your class a map of the U.S. Find the location of the two schools and ask students how to find the distance between them (use the scale on the map). Ask how well the two schools line up north-to-south. Discuss how an east-west displacement might affect the outcome of the experiment. (Convert the difference in longitudes of the two schools into a time difference, using ratio and proportion and the fact that 360° of longitude corresponds to 24 hours. Then compare the difference with the uncertainty in identifying local noon.)

Note that near local noon, the shadow length does not change much with time. For this reason, missing local noon by a few minutes is not important. Suggest that students practice measuring the shadow length at noon in advance. If you need to look up the time of local noon, see the last reference.

Ask students how to select a suitable date for both schools to make measurements. Should they check the weather forecast first?

Try making the measurement of the shadow length yourself. If possible, drive the stick into the ground, and check with the level, or a plumb line, to be sure that the stick is vertical.

Tape copy paper on top of the sheet of cardboard, and check with the level to make sure that this surface is horizontal. Be sure your stick is short enough so the shadow doesn't extend off the paper. Ask students to describe what height they will measure (the distance from the top of the stick to the surface where they see the shadow).



Demonstration

Students may have difficulty understanding how the method presented here, with two schools along the same north-south line (the same meridian) working together, permits them to measure Earth's radius on any day of the year. If you have a globe, mount two straw pieces on the same meridian. Place the globe in the beam of an overhead projector. Ask a student to rotate the globe so the location of one straw is at local noon. Ask what time it is at the other straw (local noon also). How can they tell? (The shadows have minimum length.) Have a student change the orientation of the globe axis and repeat. Ask students to relate this demonstration to the geometry of Figures 4 and 6.

A different way to visualize the geometry is to imagine the plane containing the center of Earth and points *A* and *B*. As Earth turns on its axis, this plane sweeps through all of space. When this plane is oriented so the

sun lies within it, then the shadow of each stick lies within the plane as well, so it is local noon at the location of each stick.

Assessment

There are several ways to assess students' performance in the Eratosthenes project.

- Students can choose one or more of the first three objectives and write or present the description that is specified in those objectives.
- Students can present a portfolio of student work, explanations, and drawings to show how they measured the radius of Earth.
- Students can prepare a written or oral presentation to a younger student on Eratosthenes' measurement of the circumference of Earth.

References

WYP Eratosthenes Project

<http://www.physics2005.org/events/eratosthenes/>

Center for Improved Engineering and Science Education

<http://www.k12science.org/noonday/askanexpert.html>

NASA

http://heasarc.gsfc.nasa.gov/docs/cosmic/earth_info.html

University of California, Berkeley

<http://astron.berkeley.edu/~krumholz/sq/astro/class1.txt>

U.S. Naval Observatory

Time of local noon ("sun transit")

http://aa.usno.navy.mil/data/docs/RS_OneDay.html

ERATOSTHENES 2019




Fig.1 The collaborating cities

ERATOSTHENES 2019



Fig.2 The geographical coordinates

ERATOSTHENES 2019





Fig.3 The North-South distances

ERATOSTHENES 2019

Worksheet

for

The measurement of the Radius of the Earth



North – South Distances AB (Km)			
	Amaliada	Genoa	Kassel
Amaliada			
Genoa			
Kassel			

Table 1

Measurements	Stick height	Shadow length
Amaliada	$h_1 = \dots\dots\dots$ cm	$l_1 = \dots\dots\dots$ cm
Genoa	$h_2 = \dots\dots\dots$ cm	$l_2 = \dots\dots\dots$ cm
Kassel	$h_3 = \dots\dots\dots$ cm	$l_3 = \dots\dots\dots$ cm

Table 2

Calculations			
Amaliada	$\tan \theta_1 = \frac{l_1}{h_1} = \dots\dots\dots$	$\theta_1 = \dots\dots\dots$	$\varphi = \theta_1 - \theta_i = \dots\dots\dots$
Genoa	$\tan \theta_2 = \frac{l_2}{h_2} = \dots\dots\dots$	$\theta_2 = \dots\dots\dots$	$\varphi = \theta_2 - \theta_i = \dots\dots\dots$
Kassel	$\tan \theta_3 = \frac{l_3}{h_3} = \dots\dots\dots$	$\theta_3 = \dots\dots\dots$	$\varphi = \theta_3 - \theta_i = \dots\dots\dots$

Table 3

ERATOSTHENES 2019

$$\frac{AB}{\varphi} = \frac{\text{Circumference}}{360^\circ} \Rightarrow$$

$$\text{Circumference} = \frac{AB \cdot 360^\circ}{\varphi} \Rightarrow$$

$$\text{Circumference} = \dots\dots\dots \text{Km}$$

$$\text{EarthRadius}(R_{\oplus}) = \frac{\text{Circumference}}{2\pi} \Rightarrow$$

$$R_{\oplus} = \dots\dots\dots \text{Km}$$

Eratosthenes Experiment



«The Eratosthenes experiment for the measurement of the Radius of the Earth - 2019»:

An inter-scientific and educative action!

Description

On Wednesday, March 20th 2019, more than 20.000 students and 1500 teachers at more of 650 schools in Greece, just about the time the sun was at its highest point in the sky, using simple instrumentation, followed the steps and thoughts of the great scientist Eratosthenes, as happened in 240 BC. From northern Greece to Cyprus and from Imvros to Corfu, at noon on that day, the Eratosthenes' disciples and friends of learning, managed to calculate the circumference of the Earth, in an amazing fashion and with minimal error.

Our students put sticks or other objects on the ground, measured their height and shadow's length and with simple mathematical reasoning and the aid of modern technologies (Google Earth, Sun Calc, etc.) calculated the circumference and the radius of the Earth.




Fig.1 The collaborating cities




Fig.3 The North-South distances



Fig.2 The geographical coordinates

The 2nd Junior High School of Amaliada, for fourth consecutive year, participated in this action with a group of students of the third grade.


These students, members of the Erasmus+ group of our school, exchanged their data with the students of the collaborating I.I.S.S. Einaudi-Casaregis-Galileo Galilei school in Genoa, Italy, and calculated the Earth's radius with minimal error, in a rather cloudy day.

Being part of the program ***European Schools Go Green***, our students showed that the Sun and the light it emits can not only be used as an energy source but also as a tool for the evaluation of impossible things, like the Earth's radius.

The conduction of this interdisciplinary experiment was made with the help of the teacher

Vassilopoulos Grigoris, Physicist

North – South Distances AB (Km)		
	Amaliada	Genoa
Amaliada		
Genoa	732.45	



Measurements	Stick height	Shadow length
Amaliada	$h_1 = 130$ cm	$l_1 = 102$ cm
Genoa	$h_2 = 95$ cm	$l_2 = 98$ cm

$$\frac{AB}{\varphi} = \frac{Circumference}{360^\circ} \Rightarrow$$

$$Circumference = \frac{AB \cdot 360^\circ}{\varphi} \Rightarrow$$

$$Circumference = \frac{732.45 \text{ Km} \cdot 360^\circ}{7.8^\circ} \Rightarrow$$

$$Circumference = 33805 \text{ Km}$$

$$\text{EarthRadius}(R_{\oplus}) = \frac{\text{Circumference}}{2\pi} \Rightarrow$$

$$R_{\oplus} = 5383 \text{ Km}$$


CONCLUSION

Because the experiment was done at precise the spring equinox, the angle that each school should measure, corresponds with the actual geographical latitude of each city. As it has not been possible to perfectly coincide with it, an error has occurred in the measurement of the Earth's radius.

	Latitude	Measured angle	Deviation	Measured radius	Earth radius	Error in radius
Amaliada	37.8°	38.1°	0.3°	5383 Km	6371 Km	15.5%
Genoa	44.4°	45.9°	1.5°			







2ND Junior High School
of Amaliada Greece



Galileo Galilei Technical High
School of Genova, Italy



Genova



Amaliada



Genova



Amaliada









ERASMUS PORTRAIT SUN STENCIL ART BY STUDENTS

ERATOSTHENES PORTRAIT SUN STENCIL ART BY STUDENTS



We have designed a protractor that might suit our measurements and we have built it using the 3D printer.





We have made a rehearsal the day before to test our tools.

The day we set our meeting with the Greek school was a sunny one. We have been lucky!



We made our calculations with just one grade difference. But Eratosthenes himself wasn't successful the first time!

ERATOSTHENES 2019

Worksheet
for
The measurement of the Radius of the Earth

North - South Distances AB (Km)

	Amaliada	Genoa	Kassel
Amaliada		735	
Genoa	732.45		
Kassel			

Table 1

Measurements	Stick height	Shadow length
Amaliada	$h_1 = 130$ cm	$l_1 = 102$ cm
Genoa	$h_2 = \dots 95 \dots$ cm	$l_2 = \dots 98 \dots$ cm
Kassel	$h_3 = \dots$ cm	$l_3 = \dots$ cm

Table 2

Calculations			
Amaliada	$\tan \theta_1 = \frac{l_1}{h_1} = 0.785$	$\theta_1 = 38.1^\circ$	$\varphi = \theta_1 - \theta_2 = \dots$
Genoa	$\tan \theta_2 = \frac{l_2}{h_2} = \dots 1.03 \dots$	$\theta_2 = \dots 45.9^\circ$	$\varphi = \theta_2 - \theta_1 = \dots 7.8^\circ$
Kassel	$\tan \theta_3 = \frac{l_3}{h_3} = \dots$	$\theta_3 = \dots$	$\varphi = \theta_3 - \theta_1 = \dots$

Table 3

ERATOSTHENES 2019

$$\frac{AB}{\varphi} = \frac{\text{Circumference}}{360^\circ} \Rightarrow$$

$$\text{Circumference} = \frac{AB \cdot 360^\circ}{\varphi} \Rightarrow \frac{735 \cdot 360^\circ}{7.8} = 33.923 \text{ km}$$

$$\text{Circumference} = \dots \text{Km}$$

$$\text{EarthRadius}(R_\oplus) = \frac{\text{Circumference}}{2\pi} \Rightarrow$$

$$R_\oplus = \dots \text{Km}$$

GENOVA: $44,416^\circ$ N $8,908^\circ$ E

Thanks to our teachers:

Mr. Giuseppe Canepa

Mr. Andrea Boccalero



Mr. Corrado Campi
Mr. Pietro Belmonte
Ms. Franca Monzeglio



According to Physics World magazine, the Eratosthenes experiment is one of the 10 most beautiful experiments in the history of Physics.

The list

1. Young's double slit experiment (1801)
2. Galileo's experiment on the free fall of objects (1600)
3. Millikan's experiment on the calculation of the charge of the electron (1910)
4. The analysis of sun light into spectrum by Newton (1665-1666)
5. Young's experiment on the Interference of light (1801)
6. Cavendish's torsion-bar experiment (1798)
7. Eratosthenes' measurement of the Earth's circumference (3rd. century B.C.)
8. Galileo's experiments with rolling balls down inclined planes (1600s)
9. Rutherford's discovery of the nucleus (1911)
10. Foucault's pendulum (1851)



(New York Times, 25 September 2002)

Which one of the other " most beautiful experiments in the history of Physics " would you like to conduct in collaboration with your Erasmus school partners?

VOTE IN OUR MINI SURVEY HERE:

<https://pollev.com/AANDREOU019>

Which one of the other " most beautiful experiments in the history of Physics " would you like to conduct in collaboration with your Erasmus school partners?

You can respond once

0

Young's double slit experiment (1801)

0

Galileo's experiment on the free fall of objects (1600)

0

Millikan's experiment on the calculation of the charge of the electron (1910)

0

The analysis of sun light into (samiamidi)! spectrum by Newton (1665-1666)

0

Young's experiment on the Interference of light (1801)


0

Cavendish's torsion-bar experiment (1798)


0

Galileo's experiments with rolling balls down inclined planes (1600s)

Eratosthenes experiment results 2019



The measurement of the Radius of the Earth



The meaning of the experiment

- The Junior High School of Amaliada, the Galileo Galilei Gymnasium and the Goethe High School of Kassel perform all the same experiment
- With the results we can calculate the radius of the earth

The measurement

- Every day at ca. 12:30 CET
- Accurate calculation of the sun with „suncalc.net“
- Measuring with a 1m tall piece of wood and a measuring tape



The calculation

- To get the result, the length of the shadow is divided by the length of the piece of wood

Calculations			
Amaliada	$\tan \theta_1 = \frac{l_1}{h_1} = \dots\dots\dots$	$\theta_1 = \dots\dots\dots$	$\varphi = \theta_1 - \theta_i = \dots\dots\dots$
Genoa	$\tan \theta_2 = \frac{l_2}{h_2} = \dots\dots\dots$	$\theta_2 = \dots\dots\dots$	$\varphi = \theta_2 - \theta_i = \dots\dots\dots$
Kassel	$\tan \theta_3 = \frac{l_3}{h_3} = \dots\dots\dots$	$\theta_3 = \dots\dots\dots$	$\varphi = \theta_3 - \theta_i = \dots\dots\dots$

The results

- Day 1: 0cm
- Day 2: 122,7cm
- Day 3: 119,9cm
- Day 4: 117,8cm
- Day 5: 116,4cm
- The measuring went well, disregarding from the first day.